



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 7/06, 7/48	A1	(11) International Publication Number: WO 99/24003 (43) International Publication Date: 20 May 1999 (20.05.99)
(21) International Application Number: PCT/IB98/01771 (22) International Filing Date: 6 November 1998 (06.11.98) (30) Priority Data: 08/966,834 10 November 1997 (10.11.97) US (71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors: DRZEWIECKI, Paul, Joseph; 3049 So- cialville-Foster Road, Maineville, OH 45039 (US). LISTRO, Joseph, Anthony; 9680 Waterford Place #103, Loveland, OH 45140 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217-1087 (US).		(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: COMPOSITIONS CONTAINING COMBINATIONS OF LIQUID POLYOL FATTY ACID POLYESTERS AND A LIQUID OIL (57) Abstract <p>Disclosed are compositions suitable for topical application to human hair or skin, which comprise (a) from about 0.1 % to about 99.9 % by weight of a liquid polyol fatty acid polyester having a melting point of from about -30 °C to about 30 °C, wherein the liquid polyol fatty acid polyester has a polyol moiety and at least 1 fatty acid moiety, the polyol moiety having at least 4 free hydroxyl groups wherein at least 60 % of these free hydroxyl groups are esterified with one or more fatty acid moieties having from about 8 to about 22 carbon atoms; and (b) from about 0.1 % to about 99.9 % by weight of a liquid oil having a melting point of from about -30 °C to less than about 30 °C, wherein the liquid oil is substantially free of liquid polyol fatty acid polyesters, isohexadecane, isopropyl palmitate, or blends of C₁₃-C₁₄ isoparaffins and polyacrylamide and laureth-7. The composition preferably further comprises a topical carrier for the liquid polyol fatty acid polyester and liquid oil combination. The composition provides effective emolliency and aesthetic benefits.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

COMPOSITIONS CONTAINING COMBINATIONS OF LIQUID POLYOL FATTY ACID POLYESTERS AND A LIQUID OIL

TECHNICAL FIELD

The present invention relates to compositions suitable for topical application to human hair or skin, which comprise a select combination of liquid polyol fatty acid polyesters having a melting point of from about -30°C to about 30°C and a select liquid oil having a melting point of from about -30°C to less than about 30°C.

BACKGROUND OF THE INVENTION

Topical compositions containing emollients have been used for many years in the treatment of human hair or skin. For example, occlusive hydrocarbons such as petrolatum have been used as topical emollients to provide a protective film onto human skin to prevent water loss to the environment. Petrolatum has also been used in hair care products such as conditioners and grooming aids.

However, the most effective and widely used compositions containing occlusive emollients suffer from negative aesthetic qualities such as greasiness and stickiness. Moreover, some occlusive emollients that are used to provide a protective film on the skin results in clogging the skin pores and preventing the flow of oxygen. This obstruction of the skin surface or blockage of the passage or circulation of air and moisture limits the use of such heavy, occlusive emollient materials. Additionally, European Patent No. 458,600 B1, published March 2, 1994, discloses occlusive skin care compositions containing a polyol fatty acid polyester having at least 4 free hydroxyl groups, at least 60% of which are esterified with one or more fatty acids having from 8 to 22 carbon atoms which can form an occlusive film on the skin following topical application thereon. U.S. Patent No. 5,160,738, to Macaulay et al., issued November 3, 1992, further discloses occlusive compositions containing a blend of two or more polyol fatty acid polyesters which has the appearance and physical properties of petrolatum. However, these compositions also have the disadvantage of being heavy and can clog the skin pores preventing the flow of oxygen. Therefore, the need exists for materials which can provide emolliency and acceptable aesthetic qualities without being heavy and occlusive.

It has now been found that compositions containing nonocclusive emollients can be formulated without being heavy, sticky, or greasy. These compositions contain a select combination of liquid polyol fatty acid polyesters having a melting point of from about -30°C to about 30°C, and a liquid oil having a melting point of from about -30°C to less than about 30°C, wherein the liquid oil is substantially free of liquid polyol fatty acid polyesters, isohexadecane, isopropyl palmitate, or blends of C₁₃-C₁₄ isoparaffins and polyacrylamide and laureth-7. These compositions are applicable in a variety of products to provide both emolliency and aesthetic benefits.

It is therefore an object of the present invention to provide a composition which has both effective emolliency and acceptable aesthetic qualities, and which contain a nonocclusive emollient used in

combination with a select liquid oil. It is a further object of the present invention to provide topical compositions containing polyol fatty acid polyesters which are not heavy, sticky, and greasy, and which are effective in the treatment of human hair or skin.

SUMMARY OF THE INVENTION

The present invention relates to a composition suitable for topical application to the human hair or skin, which comprises (a) from about 0.1% to about 99.9% by weight of a liquid polyol fatty acid polyester having a melting point of from about -30°C to about 30°C, wherein the liquid polyol fatty acid polyester has a polyol moiety and at least 1 fatty acid moiety, the polyol moiety having at least 4 free hydroxyl groups wherein at least 60% of these free hydroxyl groups are esterified with one or more fatty acid moieties having from about 8 to about 22 carbon atoms; and (b) from about 0.1% to about 99.9% by weight of a liquid oil having a melting point of from about -30°C to less than about 30°C, wherein said liquid oil is substantially free of liquid polyol fatty acid polyesters, isohexadecane, isopropyl palmitate, or blends of C₁₃-C₁₄ isoparaffins and polyacrylamide and laureth-7. The composition preferably further comprises a topical carrier for the liquid polyol fatty acid polyester and liquid oil combination.

All percentages and ratios used herein are by weight and all measurements made are at 25°C, unless otherwise designated. The invention hereof can comprise, consist of, or consist essentially of, the essential as well as optional ingredients and components described herein.

DETAILED DESCRIPTION OF THE INVENTION

The term "topical composition" as used herein means a composition suitable for topical application to human hair or skin. The term is used to encompass a wide variety of personal care, beauty care, and cosmetic compositions. Nonlimiting examples of topical compositions include lotions, creams, hand and body lotions, skin conditioning lotions and creams, skin protectant compositions, sunscreen compositions, cold creams, anti-acne compositions, skin renewal products, non-lathering cleansing lotions, moisturizers, facial moisturizers, make-ups, foundations, lipsticks, lip protectants, skin cleansers, hand, face, and body cleansers, shower products, shampoos, and the like.

The term "topical carrier", as used herein, is well-known to one of ordinary skill in the art, and means one or more compatible solid or liquid filler diluents or vehicles which are suitable for administration to a human. The term "compatible", as used herein, means that the components of the topical carrier are capable of being commingled with the components of the present invention, and with each other, in a manner such that there is no interaction which would substantially reduce the efficacy or aesthetics of the cosmetic composition under ordinary use situations. The topical carrier must be a pharmaceutically acceptable carrier. The term "pharmaceutically-acceptable", as used herein, means that the topical carrier must be of sufficiently high purity and be suitable for use in contact with human hair or skin without undue toxicity, incompatibility, instability, allergic response, and the like.

The term "liquid", as used herein, refers to materials which are flowable fluids under ambient conditions of one atmosphere of pressure, at about 50% relative humidity, and at from about 20°C to about 25°C, unless otherwise specified.

The term "select liquid oils", as used herein, refers to liquid oil materials which have a melting point of less than about 30°C, and which are substantially free of liquid polyol fatty acid polyesters, isohexadecane, isopropyl palmitate, or blends of C₁₃-C₁₄ isoparaffins and polyacrylamide and laureth-7. In this context, the term "substantially free" means that the compositions preferably contain less than an effective-amount of such excluded materials when used in combination with the liquid polyol fatty acid polyester to provide any emolliency or aesthetics benefits. Generally, the compositions contain less than 1%, preferably less than 0.5%, more preferably less than 0.25%, even more preferably less than 0.1%, most preferably zero percent, of such excluded materials by weight of the composition.

The melting point of the liquid polyol fatty acid polyesters and liquid oils can be determined using conventional techniques. Such techniques are well defined in the art, and include thermometric as well as calorimetry methodology. A particularly preferred technique for determining the melting point is described in U.S. Patent No. 5,306,514, to Letton et al., issued April 26, 1994, which is incorporated by reference herein in its entirety. This technique typically involves measuring melting points using Differential Scanning Calorimetry (DSC) whereby a scanning temperature of 5°C/minute is used to measure the melting point. The melting point is the temperature at the intersection of the baseline, i.e. the specific heat line, with the line tangent to the trailing edge of the endothermic peak.

Liquid Polyol Fatty Acid Polyester

The composition of the present invention comprises a nonocclusive liquid polyol fatty acid polyester at concentrations ranging from about 0.1% to about 99.9%, preferably from about 0.5% to about 75%, more preferably from about 1% to about 50%, even more preferably from about 2% to about 25%, by weight of the composition. These liquid polyol fatty acid polyesters have melting points below about 30°C, and are derived from any aliphatic or aromatic polyol which has at least 4 free hydroxyl groups, of which at least 60% of these free hydroxyl groups are then esterified with one or more fatty acids having from about 8 to about 22 carbon atoms. The fatty acids can also be described as carboxylic acids, because the terms fatty acid and carboxylic acid are often used interchangeably by those skilled in the art.

The liquid polyol polyesters employed in this invention comprise certain polyols, especially sugars or sugar alcohols, esterified with one or more fatty acid groups. Accordingly, the polyol starting material must have at least four esterifiable hydroxyl groups. Examples of preferred polyols are sugars, including monosaccharides and disaccharides, and sugar alcohols. Examples of monosaccharides containing four hydroxyl groups are xylose and arabinose and the sugar alcohol derived from xylose, which has five hydroxyl groups, i.e., xylitol. The monosaccharide, erythrose, is not suitable in the practice of this invention since it only contains three hydroxyl groups, but the sugar alcohol derived from erythrose, i.e., erythritol, contains four hydroxyl groups and accordingly can be used. Suitable five hydroxyl group-

containing monosaccharides are galactose, fructose, and sorbose. Sugar alcohols containing six hydroxyl groups derived from the hydrolysis products of sucrose, as well as glucose and sorbose, e.g., sorbitol, are also suitable. Examples of disaccharide polyols which can be used include maltose, lactose, and sucrose, all of which contain eight hydroxyl groups.

The polyols used in the liquid polyol esters of the present invention have from about 4 to about 12, more preferably from about 4 to about 11, and most preferably from about 4 to about 8 hydroxyl groups. Preferred polyols for preparing the polyesters for use in the present invention are selected from the group consisting of erythritol, xylitol, sorbitol, glucose, and sucrose. Sucrose is especially preferred.

The preferred polyol starting material having at least four hydroxyl groups must be esterified on at least 60% of the hydroxyl groups with a fatty acid containing from about 8 to about 22 carbon atoms, preferably from about 8 to about 18 carbon atoms. Examples of such fatty acids include caprylic, capric, lauric, myristic, myristoleic, palmitic, palmitoleic, stearic, oleic, ricinoleic, linoleic, linolenic, eleostearic, arachidic, arachidonic, behenic, and erucic acids. The fatty acids can be derived from naturally occurring or synthetic fatty acids; they can be saturated or unsaturated, including positional and geometrical isomers. However, in order to provide liquid polyesters of the type used herein, at least about half of the fatty acid incorporated into the polyester molecule must be unsaturated. Oleic and linoleic acids, and mixtures thereof, are especially preferred.

The liquid polyol fatty acid polyesters useful in this invention must contain one or more fatty acid ester groups. It is not necessary that all of the hydroxyl groups of the polyol be esterified with fatty acids, but it is preferable that at least 60% of the hydroxyl groups are esterified with fatty acid ester groups. Most preferably, substantially all of the hydroxyl groups of the polyol are esterified with fatty acids, i.e., the polyol moiety is substantially completely esterified. The fatty acids esterified to the polyol molecule can be the same or mixed, but as noted above, a substantial amount of the unsaturated acid ester groups must be present to provide liquidity.

To illustrate the above points, a sucrose di-fatty acid ester would be suitable for use herein, but is not preferred because it has more than two unesterified hydroxyl groups. Likewise, a sucrose tetra-fatty acid ester would be suitable, but is not preferred because it also has more than two unesterified hydroxyl groups. Highly preferred compounds in which all the hydroxyl groups are esterified with fatty acids include the liquid sucrose octa-substituted fatty acid esters.

The following are non-limiting examples of specific liquid polyol fatty acid polyesters containing one or more fatty acid ester groups suitable for use in the present invention: glucose oleate, the glucose ester of soybean oil fatty acids (unsaturated), the mannose ester of mixed soybean oil fatty acids, the galactose ester of oleic acid, the arabinose ester of linoleic acid, xylose linoleate, sorbitol oleate, sucrose oleate, glucose dioleate, the glucose diesters of soybean oil fatty acids (unsaturated), the mannose diesters of mixed soybean oil fatty acids, the galactose diesters of oleic acid, the arabinose diesters of linoleic acid, xylose dilinoleate, sorbitol dioleate, sucrose dioleate, glucose trioleate, the glucose triesters of soybean oil

fatty acids (unsaturated), the mannose triesters of mixed soybean oil fatty acids, the galactose triesters of oleic acid, the arabinose triesters of linoleic acid, xylose trilinoleate, sorbitol trioleate, sucrose trioleate, the glucose tetraesters of soybean oil fatty acids (unsaturated), the mannose tetraesters of mixed soybean oil fatty acids, the galactose tetraesters of oleic acid, the arabinose tetraesters of linoleic acid, xylose tetralinoleate, sorbitol tetraoleate, galactose pentaoleate, the sorbitol hexaesters of unsaturated soybean oil fatty acids, xylitol pentaoleate, sucrose tetraoleate, sucrose pentaoleate, sucrose hexaoleate, sucrose heptaoleate, sucrose octaoleate, and mixtures thereof. Preferred are liquid polyol esters selected from the group consisting of sucrose pentaoleate, sucrose hexaoleate, sucrose heptaoleate, sucrose octaoleate, and mixtures thereof. More preferred are sucrose hexaoleate, sucrose heptaoleate, sucrose octaoleate, and mixtures thereof.

The preferred liquid polyol fatty acid polyesters of the present invention have melting points of from about -30°C to about 30°C , preferably from about -30°C to about 27.5°C , and more preferably from about -30°C to about 25°C . The melting points are measured using conventional techniques.

The polyol fatty acid polyesters suitable for use herein can be prepared by a variety of methods well known to those skilled in the art. These methods include: transesterification of the polyol with methyl, ethyl or glycerol fatty acid esters using a variety of catalysts; acylation of the polyol with a fatty acid chloride; acylation of the polyol with a fatty acid anhydride; and acylation of the polyol with a fatty acid, per se. See U.S. Patent No. 2,831,854; U.S. Patent No. 4,005,196, to Jandacek, issued January 25, 1977; and U.S. Patent No. 4,005,196, to Jandacek, issued January 25, 1977, all of which are incorporated by reference herein in their entirety.

Liquid Oil

The composition of the present invention comprises a select liquid oil, as defined herein, used in combination with the liquid polyol fatty acid polyester described herein. Concentrations of the select liquid oil range from about 0.1% to about 99.9%, preferably from about 0.5% to about 75%, more preferably from about 1% to about 50%, even more preferably from about 2% to about 25%, by weight of the composition.

Liquid oils useful herein are those materials which are substantially free of liquid polyol fatty acid polyesters, isohexadecane, isopropyl palmitate, or blends of C₁₃-C₁₄ isoparaffins and polyacrylamide and laureth-7, and which have a melting point of less than about 30°C, preferably from about -30°C to less than about 30°C, more preferably from about -30°C to about 27.5°C, and even more preferably from about -30°C to about 25°C. The liquid oils generally have a low solubility in water, generally less than about 1% by weight at 25°C. Examples of suitable liquid oils include, but are not limited to, mineral oil, hydrocarbons having from about 5 to about 16 carbon atoms, fatty alcohol esters having from about 3 to about 22 carbon atoms, fatty acid esters having from about 3 to about 30 carbon atoms, vegetable oils, and mixtures thereof. Other liquid oils suitable for use herein are described in WO 95-00166, to Gordon et al., published January 5, 1995, which is incorporated by reference herein in its entirety.

Mineral oil, which is also known as petrolatum liquid, is suitable for use herein. This is a mixture of liquid hydrocarbons obtained from petroleum. See The Merck Index, Tenth Edition, Entry 7048, p. 1033 (1983) and International Cosmetic Ingredient Dictionary, Fifth Edition, vol. 1, p.415-417 (1993), which are incorporated by reference herein in their entirety.

Examples of suitable select hydrocarbon liquid oils for use herein include straight and branched chain hydrocarbons having from about 5 to about 16 carbon atoms. Nonlimiting examples of these hydrocarbon materials include dodecane, isododecane, hydrogenated polyisobutylene, and hexadecane. Also useful are the C₅-C₁₆ isoparaffins, excluding the C₁₃-C₁₄ isoparaffins when these compounds are used in combination with polyacrylamide and laureth-7. These C₅-C₁₆ isoparaffins are also referred to as C₅-C₁₆ branched hydrocarbons.

Fatty alcohol esters suitable for use as a select liquid oil defined herein include esters and diesters of fatty alcohols which have from about 3 to about 22 carbon atoms. Also useful are fatty acid esters having from about 3 to about 28 carbon atoms. Nonlimiting examples of these ester materials include isopropyl myristate, isopropyl stearate, diisopropyl adipate, and dioctyl sebacate (dioctyl ester of decanedioic acid).

Other suitable select liquid oils for use herein include vegetable oils which are liquid at ambient temperatures of from about 20°C to about 25°C. Examples of suitable vegetable oils include cod-liver oil, dolphin oil, lard oil, neat's-foot oil, porpoise oil, seal oil, sperm oil, whale oil, acorn oil, almond oil, beechnut oil, chaulmoogra oil, rapeseed, soybean oil, sunflower-seed oil, groundnut oil, cottonseed oil, corn oil, safflower oil, olive oil, menhaden oil, sesame seed, castor oil, hazelnut oil, hemp-seed oil, linseed

oil, mustard (black) oil, neem oil, peanut oil, pistachio-nut oil, poppy-seed oil, pumpkin-seed oil, Tung oil, white-mustard-seed oil, pine oil, wheat germ oil, derivatives thereof and mixtures thereof.

Topical Carrier

The composition of the present invention comprises from about 0.1% to about 99.9%, preferably from about 50% to about 99%, and more preferably from about 60% to about 95% by weight of a topical carrier for the liquid polyol fatty acid polyester and liquid oil combination, and for any other optional components of the present invention.

~~The liquid-polyol-fatty-acid-polyester-and-liquid-oil-combination-of-the-present-invention-can-be~~ formulated into a wide variety of product types, including creams, lotions, milks, gels, hand and body lotions, cold creams, non-lathering cleansing lotions, facial moisturizers, sunscreens, anti-acne preparations, topical analgesics, mascaras, lipsticks, skin cleansers, hand, face, and body cleansers, shower products, shampoos, and the like. The carriers and any additional components required to formulate such products vary with product type and can be routinely chosen by one skilled in the art.

The topical carrier can be in a wide variety of forms. For example, emulsion carriers, including, but not limited to, oil-in-water, water-in-oil, water-in-oil-in-water, and oil-in-water-in-silicone emulsions, are useful herein. These emulsions can cover a broad range of viscosities, e.g., from about 100 cps to about 200,000 cps. Other suitable topical carriers include anhydrous liquid solvents such as alcohols, and silicones (e.g., ethanol, isopropanol, dimethicone, cyclomethicone, and the like); aqueous-based single phase liquid solvents (e.g., hydro-alcoholic solvent systems); and thickened versions of these anhydrous and aqueous-based single phase solvents (e.g., where the viscosity of the solvent has been increased to form a solid or semi-solid by the addition of appropriate gums, resins, waxes, polymers, salts, and the like). Examples of topical carrier systems useful in the present invention are described in the following references all of which are incorporated herein by reference in their entirety: "Sun Products Formulary" Cosmetics & Toiletries, vol. 105, pp. 122-139 (December 1990); "Sun Products Formulary", Cosmetics & Toiletries, vol. 102, pp. 117-136 (March 1987); U.S. Patent No. 4,960,764 to Figueroa et al., issued October 2, 1990; U.S. Patent No. 4,254,105 to Fukuda et al., issued March 3, 1981; U.S. Patent No. 4,976,953, to Orr et al., issued December 11, 1990; U.S. Patent No. 5,073,372, to Turner et al., issued December 17, 1991; U.S. Patent No. 5,585,104, to Ha et al., issued December 17, 1996; U.S. Patent No. 5,607,678, to Moore et al., issued March 4, 1997; U.S. Patent No. 5,607,980, to McAtee et al., issued March 4, 1997; and U.S. Patent No. 5,618,522, to Kaleta et al., issued April 8, 1997.

The topical carrier can also comprise an oil-in-water emulsion system having complex structures such as liquid crystals and crystalline gel networks. The nature of liquid crystals, the formation of liquid crystals, the properties and advantages of liquid crystals are described further in G. Dahms, Properties of O/W Emulsions With Anisotropic Lamellar Phases, 101 *Cosmetics & Toiletries*, 113-115 (1986); P. Loll, Liquid Crystals in Cosmetic Emulsions, *ICI Surfactants' Publication RP94-93E*; and G. M. Eccleston,

Multiple-Phase Oil-In-Water Emulsions, 41, *J. Soc. Cosmet. Chem.*, 1-22, (January/February 1990); all of which are incorporated herein by reference in their entirety.

Additional Components

A wide variety of additional components can be incorporated into the compositions herein. Non-limiting examples include the following:

Pharmaceutical Actives

The compositions of the present invention can comprise a safe and effective amount of a pharmaceutical active. The phrase "safe and effective amount", as used herein, means an amount of an active high enough to significantly or positively modify the condition to be treated, but low enough to avoid serious side effects (at a reasonable benefit/risk ratio), within the scope of sound medical judgment. A safe and effective amount of the pharmaceutical active will vary with the specific active, the ability of the composition to penetrate the active through the skin, the amount of composition to be applied, the particular condition being treated, the age and physical condition of the patient being treated, the severity of the condition, the duration of the treatment, the nature of concurrent therapy, and like factors.

The pharmaceutical actives which can be used in the compositions of the present invention preferably comprise from about 0.1% to about 20% by weight of the compositions, more preferably from about 0.1% to about 10%, and most preferably from about 0.1% to about 5%. Mixtures of pharmaceutical actives may also be used.

Nonlimiting examples of pharmaceutical actives can include the following:

Useful pharmaceutical actives in the compositions of the present invention include anti-acne drugs. Anti-acne drugs for use herein include the keratolytics such as salicylic acid, sulfur, lactic acid, glycolic, pyruvic acid, resorcinol, and N-acetylcysteine; retinoids such as retinoic acid and its derivatives (e.g., cis and trans); antibiotics and antimicrobials such as benzoyl peroxide, octopirox, erythromycin, zinc, tetracyclin, triclosan, azelaic acid and its derivatives, phenoxy ethanol and phenoxy propanol, ethylacetate, clindamycin and meclocycline; sebostats such as flavinoids; alpha and beta hydroxy acids; and bile salts such as scymnol sulfate and its derivatives, deoxycholate, and cholate. Preferred anti-acne actives are those selected from the group consisting of salicylic acid, sulfur, resorcinol, lactic acid, zinc, erythromycin, benzoyl peroxide, and mixtures thereof. More preferred is salicylic acid.

Useful pharmaceutical actives in the compositions of the present invention include non-steroidal anti-inflammatory drugs (NSAIDS). The NSAIDS can be selected from the following categories: propionic acid derivatives; acetic acid derivatives; fenamic acid derivatives; biphenylcarboxylic acid derivatives; and oxicams. All of these NSAIDS are fully described in the U.S. Patent 4,985,459 to Sunshine et al., issued January 15, 1991, incorporated by reference herein. Most preferred are the propionic NSAIDS including but not limited to aspirin, acetaminophen, ibuprofen, naproxen, benoxaprofen, flurbiprofen, fenoprofen, fenbufen, ketoprofen, indoprofen, piroprofen, carprofen,

oxaprozin, pranoprofen, mioprofen, tioxaprofen, suprofen, alminoprofen, tiaprofenic acid, fluprofen and bucloxic acid. Also useful are the steroidal anti-inflammatory drugs including hydrocortisone and the like.

Useful pharmaceutical actives in the compositions of the present invention include antipruritic drugs. Antipruritic drugs preferred for inclusion herein include pharmaceutically-acceptable salts of methdilazine and trimeprazine.

Useful pharmaceutical actives in the compositions of the present invention include anesthetic drugs. Anesthetic drugs preferred for inclusion herein include pharmaceutically-acceptable salts of ~~lidocaine, bupivacaine, chlorprocaine, dibucaine, etidocaine, mepivacaine, tetracaine, dyclonine, hexylcaine, procaine, cocaine, ketamine, pramoxine and phenol.~~

Useful pharmaceutical actives in the compositions of the present invention include antimicrobial drugs (antibacterial, antifungal, antipprotozoal and antiviral drugs). Antimicrobial drugs preferred for inclusion herein include pharmaceutically-acceptable salts of β -lactam drugs, quinolone drugs, ciprofloxacin, norfloxacin, tetracycline, erythromycin, amikacin, triclosan, doxycycline, capreomycin, chlorhexidine, chlortetracycline, oxytetracycline, clindamycin, ethambutol, metronidazole, pentamidine, gentamicin, kanamycin, lineomycin, methacycline, methenamine, minocycline, neomycin, netilmicin, paromomycin, streptomycin, tobramycin, miconazole and amantadine. Antimicrobial drugs preferred for inclusion herein include tetracycline hydrochloride, erythromycin estolate, erythromycin stearate (salt), amikacin sulfate, doxycycline hydrochloride, capreomycin sulfate, chlorhexidine gluconate, chlorhexidine hydrochloride, chlortetracycline hydrochloride, oxytetracycline hydrochloride, clindamycin hydrochloride, ethambutol hydrochloride, metronidazole hydrochloride, pentamidine hydrochloride, gentamicin sulfate, kanamycin sulfate, lineomycin hydrochloride, methacycline hydrochloride, methenamine hippurate, methenamine mandelate, minocycline hydrochloride, neomycin sulfate, netilmicin sulfate, paromomycin sulfate, streptomycin sulfate, tobramycin sulfate, miconazole hydrochloride, amantadine hydrochloride, amantadine sulfate, triclosan, octopirox, parachlorometa xyleneol, nystatin, tolnaftate and clotrimazole.

Also useful herein are sunscreens agents. A wide variety of sunscreens agents are described in U.S. Patent No. 5,087,445, to Haffey et al., issued February 11, 1992; U.S. Patent No. 5,073,372, to Turner et al., issued December 17, 1991; U.S. Patent No. 5,073,371, to Turner et al. issued December 17, 1991; and Segarin, et al., at Chapter VIII, pages 189 et seq., of Cosmetics Science and Technology, all of which are incorporated herein by reference in their entirety. Preferred among those sunscreens which are useful in the compositions of the instant invention are those selected from the group consisting of 2-ethylhexyl p-methoxycinnamate, 2-ethylhexyl N,N-dimethyl-p-aminobenzoate, p-aminobenzoic acid, 2-phenylbenzimidazole-5-sulfonic acid, octocrylene, oxybenzone, homomenthyl salicylate, octyl salicylate, 4,4'-methoxy-t-butylidibenzoylmethane, 4-isopropyl dibenzoylmethane, 3-benzylidene camphor, 3-(4-methylbenzylidene) camphor, titanium dioxide, zinc oxide, silica, iron oxide, and mixtures thereof.

Still other useful sunscreens are those disclosed in U.S. Patent No. 4,937,370, to Sabatelli, issued June 26, 1990; and U.S. Patent No. 4,999,186, to Sabatelli et al., issued March 12, 1991; these two

references are incorporated by reference herein in their entirety. The sunscreens disclosed therein have, in a single molecule, two distinct chromophore moieties which exhibit different ultra-violet radiation absorption spectra. One of the chromophore moieties absorbs predominantly in the UVB radiation range and the other absorbs strongly in the UVA radiation range. These sunscreens provide higher efficacy, broader UV absorption, lower skin penetration and longer lasting efficacy relative to conventional sunscreens. Especially preferred examples of these sunscreens include those selected from the group consisting of 4-N,N-(2-ethylhexyl)methylaminobenzoic acid ester of 2,4-dihydroxybenzophenone, 4-N,N-(2-ethylhexyl)methylaminobenzoic acid ester with 4-hydroxydibenzoylmethane, 4-N,N-(2-ethylhexyl)methylaminobenzoic acid ester of 2-hydroxy-4-(2-hydroxyethoxy)benzophenone, 4-N,N-(2-ethylhexyl)methylaminobenzoic acid ester of 4-(2-hydroxyethoxy)dibenzoylmethane, and mixtures thereof.

Generally, the compositions can comprise from about 0.5% to about 20% of the sunscreens useful herein. Exact amounts will vary depending upon the sunscreen chosen and the desired Sun Protection Factor (SPF). SPF is a commonly used measure of photoprotection of a sunscreen against erythema. See Federal Register, Vol. 43, No. 166, pp. 38206-38269, August 25, 1978, which is incorporated herein by reference in its entirety.

Also useful in the compositions of the present invention are sunless tanning agents including dihydroxyacetone, glyceraldehyde, indoles and their derivatives, and the like. These sunless tanning agents can also be used in combination with the sunscreen agents.

Other useful actives include skin bleaching (or lightening) agents including but not limited to hydroquinone, ascorbic acid, kojic acid and sodium metabisulfite.

Humectants and Moisturizers

The compositions of the present invention can also contain one or more additional humectant or moisturizing materials other than those described herein. A variety of these materials can be employed and each can be present at a level of from about 0.1% to about 20%, more preferably from about 1% to about 10% and most preferably from about 2% to about 5%, by weight of the composition. These materials include guanidine; glycolic acid and glycolate salts (e.g. ammonium and quaternary alkyl ammonium); lactic acid and lactate salts (e.g. ammonium and quaternary alkyl ammonium); aloe vera in any of its variety of forms (e.g., aloe vera gel); polyhydroxy alcohols such as sorbitol, glycerol, hexanetriol, propylene glycol, butylene glycol, hexylene glycol and the like; sugars and starches; sugar and starch derivatives (e.g., alkoxylated glucose); hyaluronic acid; lactamide monoethanolamine; acetamide monoethanolamine; and mixtures thereof.

Emulsifiers

The compositions herein can contain various emulsifiers. These emulsifiers are useful for emulsifying the various carrier components of the compositions herein. Suitable emulsifiers can include any of a wide variety of nonionic, cationic, anionic, and zwitterionic emulsifiers disclosed in the prior

patents and other references. See McCutcheon's, Detergents and Emulsifiers, North American Edition (1986), published by Allured Publishing Corporation; U.S. Patent No. 5,011,681 to Ciotti et al., issued April 30, 1991; U.S. Patent No. 4,421,769 to Dixon et al., issued December 20, 1983; and U.S. Patent No. 3,755,560 to Dickert et al., issued August 28, 1973; these four references are incorporated herein by reference in their entirety.

Suitable emulsifier types include esters of glycerin, esters of propylene glycol, fatty acid esters of polyethylene glycol, fatty acid esters of polypropylene glycol, esters of sorbitol, esters of sorbitan anhydrides, carboxylic acid copolymers, esters and ethers of glucose, ethoxylated ethers, ethoxylated alcohols, alkyl phosphates, polyoxyethylene fatty ether phosphates, fatty acid amides, acyl lactylates, soaps and mixtures thereof.

Suitable emulsifiers can include, but are not limited to, polyethylene glycol 20 sorbitan monolaurate (Polysorbate 20), polyethylene glycol 5 soya sterol, Steareth-20, Cetareth-20, PPG-2 methyl glucose ether distearate, Ceteth-10, Polysorbate 80, cetyl phosphate, potassium cetyl phosphate, diethanolamine cetyl phosphate, Polysorbate 60, glyceryl stearate, PEG-100 stearate, and mixtures thereof.

The emulsifiers can be used individually or as a mixture of two or more and can be included at concentrations ranging from about 0.1% to about 10%, more preferably from about 1% to about 7%, and most preferably from about 1% to about 5%, by weight of the composition.

Carboxylic Acid Copolymer Thickeners

Another component useful in the compositions herein is a carboxylic acid copolymer thickener. These crosslinked polymers contain one or more monomers derived from acrylic acid, substituted acrylic acids, and salts and esters of these acrylic acids and the substituted acrylic acids, wherein the crosslinking agent contains two or more carbon-carbon double bonds and is derived from a polyhydric alcohol. The preferred polymers for use herein are of two general types. The first type of polymer is a crosslinked homopolymer of an acrylic acid monomer or derivative thereof (e.g., wherein the acrylic acid has substituents on the two and three carbon positions independently selected from the group consisting of C₁₋₄ alkyl, -CN, -COOH, and mixtures thereof). The second type of polymer is a crosslinked copolymer having a first monomer selected from the group consisting of an acrylic acid monomer or derivative thereof (as just described in the previous sentence), a short chain alcohol (i.e. a C₁₋₄) acrylate ester monomer or derivative thereof (e.g., wherein the acrylic acid portion of the ester has substituents on the two and three carbon positions independently selected from the group consisting of C₁₋₄ alkyl, -CN, -COOH, and mixtures thereof), and mixtures thereof; and a second monomer which is a long chain alcohol (i.e. C₈₋₄₀) acrylate ester monomer or derivative thereof (e.g., wherein the acrylic acid portion of the ester has substituents on the two and three carbon positions independently selected from the group consisting of C₁₋₄ alkyl, -CN, -COOH, and mixtures thereof). Combinations of these two types of polymers are also useful herein.

In the first type of crosslinked homopolymers the monomers are preferably selected from the group consisting of acrylic acid, methacrylic acid, ethacrylic acid, and mixtures thereof, with acrylic acid being most preferred. In the second type of crosslinked copolymers the acrylic acid monomer or derivative thereof is preferably selected from the group consisting of acrylic acid, methacrylic acid, ethacrylic acid, and mixtures thereof, with acrylic acid, methacrylic acid, and mixtures thereof being most preferred. The short chain alcohol acrylate ester monomer or derivative thereof is preferably selected from the group consisting of C₁₋₄ alcohol acrylate esters, C₁₋₄ alcohol methacrylate esters, C₁₋₄ alcohol ethacrylate esters, and mixtures thereof, with the C₁₋₄ alcohol acrylate esters, C₁₋₄ alcohol methacrylate esters, and mixtures thereof, being most preferred. The long chain alcohol acrylate ester monomer is selected from C₈₋₄₀ alkyl acrylate esters, with C₁₀₋₃₀ alkyl acrylate esters being preferred.

The crosslinking agent in both of these types of polymers is a polyalkenyl polyether of a polyhydric alcohol containing more than one alkenyl ether group per molecule, wherein the parent polyhydric alcohol contains at least 3 carbon atoms and at least 3 hydroxyl groups. Preferred crosslinkers are those selected from the group consisting of allyl ethers of sucrose and allyl ethers of pentaerythritol, and mixtures thereof. These polymers useful in the present invention are more fully described in U.S. Patent No. 5,087,445, to Haffey et al., issued February 11, 1992; U.S. Patent No. 4,509,949, to Huang et al., issued April 5, 1985; U.S. Patent No. 2,798,053, to Brown, issued July 2, 1957; which are incorporated by reference herein. See also, CTFA International Cosmetic Ingredient Dictionary, fourth edition, 1991, pp. 12 and 80; which is also incorporated herein by reference.

Examples of commercially available homopolymers of the first type useful herein include the carbomers, which are homopolymers of acrylic acid crosslinked with allyl ethers of sucrose or pentaerythritol. The carbomers are available as the Carbopol® 900 series from B.F. Goodrich. Examples of commercially available copolymers of the second type useful herein include copolymers of C₁₀₋₃₀ alkyl acrylates with one or more monomers of acrylic acid, methacrylic acid, or one of their short chain (i.e. C₁₋₄ alcohol) esters, wherein the crosslinking agent is an allyl ether of sucrose or pentaerythritol. These copolymers are known as acrylates/C₁₀₋₃₀ alkyl acrylate crosspolymers and are commercially available as Carbopol® 1342, Pemulen TR-1, and Pemulen TR-2, from B.F. Goodrich. In other words, examples of carboxylic acid polymer thickeners useful herein are those selected from the group consisting of carbomers, acrylates/C_{10-C30} alkyl acrylate crosspolymers, and mixtures thereof.

The compositions of the present invention can comprise from about 0.025% to about 1%, more preferably from about 0.05% to about 0.75%, and most preferably from about 0.10% to about 0.50%, by weight of the carboxylic acid polymer thickeners.

Other Additional Components

The compositions of the present invention can comprise a wide range of other additional components. The CTFA Cosmetic Ingredient Handbook, Second Edition, 1992, which is incorporated by reference herein in its entirety, describes a wide variety of nonlimiting cosmetic and pharmaceutical

ingredients commonly used in hair and skin care industries, which are suitable for use in the compositions of the present invention. Nonlimiting examples of functional classes of ingredients are described at page 537 of this reference. Examples of these functional classes include: absorbents, abrasives, anti-acne agents, anticaking agents, antifoaming agents, antimicrobial agents, antioxidants, binders, biological additives, buffering agents, bulking agents, chelating agents, chemical additives, colorants, cosmetic astringents, cosmetic biocides, denaturants, drug astringents, film formers, fragrance components, opacifying agents, pH adjusters, plasticizers, preservatives, propellants, reducing agents, additional skin-conditioning agents, ~~suspending agents (nonsurfactant), ultraviolet light absorbers, and viscosity~~ increasing agents (aqueous and nonaqueous). Examples of other functional classes of materials useful herein that are well known to one of ordinary skill in the art include solubilizing agents, sequestrants, and the like.

Nonlimiting examples of these additional components cited in the CTFA Cosmetic Ingredient Handbook, as well as other materials useful herein, include the following: vitamins and derivatives thereof [e.g., vitamin C, Vitamin A (i.e. retinoic acid), retinol, retinoids, and the like]; anti-oxidants; polyethyleneglycols; polymers for aiding the film-forming properties and substantivity of the composition (such as a copolymer of eicosene and vinyl pyrrolidone, an example of which is available from GAF Chemical Corporation as Ganex® V-220); preservatives for maintaining the antimicrobial integrity of the compositions; antioxidants; chelators and sequestrants; crosslinked and noncrosslinked cationic polyacrylamides [e.g., Salcare SC92 which has the CTFA designation polyquaternium 32 (and) mineral oil, and Salcare SC95 which has the CTFA designation polyquaternium 37 (and) mineral oil (and) PPG-1 trideceth-6]; and aesthetic components such as fragrances, pigments, colorings, essential oils, skin senates, astringents, skin soothing agents, skin healing agents and the like, nonlimiting examples of these aesthetic components include clove oil, menthol, camphor, eucalyptus oil, eugenol, menthyl lactate, witch hazel distillate, bisabolol, dipotassium glycyrrhizinate, and the like.

Method of Use

The compositions of the present invention are used in conventional ways to provide cosmetic or pharmaceutical benefits appropriate to the product such as sun protection, anti-acne benefits, anti-wrinkle and anti-skin aging benefits, artificial tanning, analgesic benefits, skin conditioning benefits, facial moisturization, lip protection, skin cleansing benefits, grooming aids, and the like. Such methods of use depend upon the type of composition employed but generally involve topical application of an effective amount of the product to the hair or skin. By "effective amount" is meant an amount sufficient to provide the benefit desired. Typical amounts of the compositions of the present invention which are applied to the hair or skin will vary depending upon the type of composition and the benefit desired, however, typical ranges are generally from about 1 gram to about 25 gram, with about 2 gram being typical.

Examples

The following examples further describe and demonstrate embodiments within the scope of the present invention. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the present invention, as many variations thereof are possible without departing from the spirit and scope of the invention.

Ingredients are identified by chemical or CTFA name.

Example 1

A topical composition in the form of a shower product is prepared by combining the following ingredients using conventional mixing techniques. This composition utilizes a mixture of a nonocclusive liquid polyol fatty acid polyester (liquid sucrose polyester) and a liquid oil to provide emolliency and aesthetic benefits without leaving the skin feeling heavy, sticky or greasy.

<u>Ingredients</u>	<u>Weight Percent</u>
Sodium C12/14 Alkyl Ether Glycerol Sulfonate	12.0
Ammonium Laureth-3 Sulfate	3.00
Myristic Acid	1.00
Myristic Alcohol	1.00
Cocamidopropyl Betaine	3.00
Liquid Sucrose Polyester ¹	15.3
Tetrasodium EDTA	0.13
Glycerin	6.24
Perfume	0.80
Polyquat-10 (JR-30M)	0.30
Glydant	0.20
Maleated Soybean Oil	1.00
Water	QS 100

¹ Liquid sucrose polyester is a mixture of hexa-, hepta-, and octa-sucrose esters, predominately the octa-ester esterified with mixed soybean oil fatty acids.

In a suitable vessel, the Polyquat-10 is added to distilled water and allowed to mix until fully hydrated. The surfactants, and water soluble ingredients, are added and the mixture is heated with stirring to 70-80°C. In a separate vessel, the liquid sucrose polyester is combined with the liquid oil, heated with mixing until uniform, and then added to the mixture heated at 70-80°C. The mixture is allowed to cool to 25-35°C while continuing to stir. The glydant and perfume ingredients are then added with stirring, and the mixture is cooled to room temperature.

Example 2

A topical composition in the form of a shower product is prepared by combining the following ingredients using conventional mixing techniques. This composition utilizes a mixture of a nonocclusive liquid polyol fatty acid polyester (liquid sucrose polyester) and a liquid oil to provide emolliency and aesthetic benefits without leaving the skin feeling heavy, sticky or greasy.

Ingredients

Weight Percent

Ammonium Lauryl Sulfate	3.15
Ammonium Laureth-3 Sulfate	9.45
Na Lauroamphoacetate	5.40
Polyquaternium-10	0.30
Liquid Sucrose Polyester ¹	15.3
Mineral Oil	3.00
Tetrasodium EDTA	0.13
Glycerin	3.00
Perfume	0.80
Citric Acid	0.76
Lauryl Alcohol	2.00
Water	QS 100

¹ Liquid sucrose polyester is a mixture of hexa-, hepta-, and octa-sucrose esters, predominately the octa-ester esterified with mixed soybean oil fatty acids.

In a suitable vessel, the Polyquaternium-10 is added to distilled water and allowed to mix until fully hydrated. The surfactants, and water soluble ingredients, are added and the mixture is heated with stirring to 70-80°C. In a separate vessel, the liquid sucrose polyester is combined with the liquid oil and lauryl alcohol, heated with mixing until uniform, and then added to the mixture heated at 70-80°C. The mixture is allowed to cool to 25-35°C while continuing to stir. The glydant and perfume ingredients are then added with stirring, and the mixture is cooled to room temperature.

Example 3

A topical composition in the form of a moisturizer is prepared by combining the following ingredients using conventional mixing techniques. This composition utilizes a mixture of a nonocclusive liquid polyol fatty acid polyester (liquid sucrose polyester) and a liquid oil to provide emolliency and aesthetic benefits without leaving the skin feeling heavy, sticky or greasy.

<u>Ingredients</u>	<u>Weight Percent</u>
Cetyl Alcohol	1.80
Stearic Acid	0.25
Stearyl Alcohol	1.20
Peg 100-stearate	0.25
Mineral Oil	2.00
Liquid Sucrose Polyester ¹	4.00
Dimethicone 350 ²	0.50
Propyl Paraben	0.10
Arlatone (RTM) 2121 ³	1.00
Glycerin	9.00
Urea	2.00
Octyl Methoxycinnamate	2.00
Phenoxyethanol	0.25
Carbomer 1382 ⁴	0.05
Carbomer 954 ⁵	0.35
Tetrasodium EDTA	0.10
Titanium Dioxide	0.15
Methyl Paraben	0.20
NaOH	0.22

Dimethicone Q-21403 ⁶	1.00
Water	qs100

¹ Liquid sucrose polyester is a mixture of hexa-, hepta-, and octa-sucrose esters, predominately the octa-ester esterified with mixed soybean oil fatty acids.

² Dow Corning[®] 200 Fluid (350 centistoke) from Dow Corning.

³ 95% by weight sorbitan stearate and 5% by weight sucrose cocoate.

⁴ Carbopol[®] 1382 from B.F. Goodrich.

⁵ Carbopol[®] 954 from B.F. Goodrich.

⁶ Dow Corning[®] Q-2 1403 from Dow Corning which is a mixture of 85% by weight dimethicone and 15% by weight dimethiconal.

A first premix of the liquid sucrose polyester, Arlatone 2121 and other water soluble ingredients is prepared by admixing in water and heating. A second premix of oil phase ingredients other than the silicones is prepared by mixing and heating and is added to the aqueous premix. The resulting mixture is cooled. The silicones are then added to the resulting oil-in-water emulsion and the mixture is cooled before adding minor ingredients.

What is claimed is:

1. A composition comprising:
 - (a) from 0.1% to 99.9% by weight of a liquid polyol fatty acid polyester having a melting point of from -30°C to 30°C, wherein the liquid polyol fatty acid polyester has a polyol moiety and at least 1 fatty acid moiety, the polyol moiety having at least 4 free hydroxyl groups wherein at least 60% of these free hydroxyl groups are esterified with one or more fatty acid moieties having from 8 to 22 carbon atoms; and
 - (b) from 0.1% to 99.9% by weight of a liquid oil having a melting point of from -30°C to less than 30 °C, wherein the liquid oil is substantially free of liquid polyol fatty acid polyesters, isohexadecane, isopropyl palmitate, blends of C₁₃-C₁₄ isoparaffins and polyacrylamide and laureth-7, or combinations thereof.
2. A topical composition comprising:
 - (a) from 0.1% to 99.9% by weight of a liquid polyol fatty acid polyester having a melting point of from -30°C to 30°C, wherein the liquid polyol fatty acid polyester has a polyol moiety and at least 1 fatty acid moiety, the polyol moiety having at least 4 free hydroxyl groups wherein at least 60% of these free hydroxyl groups are esterified with one or more fatty acid moieties having from 8 to 22 carbon atoms;
 - (b) from 0.1% to 99.9% by weight of a liquid oil having a melting point of from -30°C to less than 30 °C, wherein the liquid oil is substantially free of liquid polyol fatty acid polyesters, isohexadecane, isopropyl palmitate, blends of C₁₃-C₁₄ isoparaffins and polyacrylamide and laureth-7, or combinations thereof; and
 - (c) from 0.1% to 99.9% by weight of a topical carrier.
3. The composition of Claim 1 or 2 wherein the fatty acid moiety contains from 8 to 18 carbon atoms.
4. The composition of any one of the preceding claims wherein the polyol moiety is selected from the group consisting of erythritol, xylitol, sorbitol, glucose, sucrose, and mixtures thereof.
5. The composition of any one of the preceding claims wherein the polyol moiety is sucrose.
6. The composition of any one of the preceding claims wherein the liquid oil is selected from the group consisting of mineral oil, hydrocarbons having from 5 to 16 carbon atoms, fatty alcohol esters having from 3 to 22 carbon atoms, fatty acid esters having from 3 to 30 carbon atoms, vegetable oils, and mixtures thereof.

7. The composition of any one of Claims 2-6 wherein the topical carrier is an oil-in-water emulsion.
8. The composition of any one of Claims 2-6 wherein the topical carrier is an anhydrous liquid solvent.
9. The composition of any one of Claims 2-8 wherein the composition is in the form of a hand lotion, body lotion, skin conditioning cream, skin protectant, sunscreen, cold cream, anti-acne composition, skin renewal product, non-lathering cleansing lotion, moisturizer, facial moisturizer, make-up, foundation, lipstick, lip protectant, hand cleanser, facial cleanser, body cleanser, shower product, shampoo, and mixtures thereof.
10. A method of treating human hair or skin comprising topically applying to a human in need of treatment a safe and effective amount of the composition of any one of the preceding claims.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 98/01771

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K7/06 A61K7/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 458 600 A (UNILEVER) 27 November 1991 see page 3, line 24 - page 4, column 27; claims 1,5-8; table 2	1-7,9,10
X	EP 0 519 727 A (UNILEVER) 23 December 1992 see page 5, line 21 - page 6, line 17; claims 1,4-6,9; table 2	1-7,9,10

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

26 January 1999

Date of mailing of the international search report

04/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Beyss, E

PCT/IB 98/01771

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 458600	A	27-11-1991	AU 6346294 A	28-07-1994
			AU 7724691 A	28-11-1991
			CA 2042980 A	25-11-1991
			DE 69101263 D	07-04-1994
			DE 69101263 T	09-06-1994
			DK 458600 T	27-06-1994
			ES 2062684 T	16-12-1994
			JP 4235114 A	24-08-1992
EP 519727	A	23-12-1992	AU 1829992 A	24-12-1992
			CA 2071365 A	22-12-1992
			JP 5194138 A	03-08-1993
			ZA 9204533 A	20-12-1993



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 7/48	A1	(11) International Publication Number: WO 00/69404 (43) International Publication Date: 23 November 2000 (23.11.00)
<p>(21) International Application Number: PCT/US00/12155</p> <p>(22) International Filing Date: 4 May 2000 (04.05.00)</p> <p>(30) Priority Data: 60/134,397 17 May 1999 (17.05.99) US 09/439,438 15 November 1999 (15.11.99) US</p> <p>(71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US).</p> <p>(72) Inventors: ZHUANG, John, Chengfeng; 464 Hilltop Lane, Cincinnati, OH 45215 (US). OBLONG, John, Erich; 6763 Little River Lane, Loveland, OH 45140 (US). BISSETT, Donald, Lynn; 3925 Dust Commander Drive, Hamilton, OH 45011 (US).</p> <p>(74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217-1087 (US).</p>		<p>(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: METHODS FOR REGULATING THE CONDITION OF MAMMALIAN KERATINOUS TISSUE VIA TOPICAL APPLICATION OF PHYTOSTEROL COMPOSITIONS</p>		
<p>(57) Abstract</p> <p>The present invention relates to methods for regulating the condition of mammalian keratinous tissue wherein the methods comprise the step of topically applying to the keratinous tissue of a mammal needing such treatment, a safe and effective amount of a composition comprising: a) a safe and effective amount of one or more phytosterols selected from the group consisting of β-sitosterol, campesterol, brassicasterol, $\Delta 5$-avenasterol, lupenol, α-spinasterol, stigmasterol, their derivatives, and combinations thereof; and b) a dermatologically acceptable carrier for the phytosterol. In additional embodiments, the above composition is suitable for thickening skin and preventing, retarding, and/or treating atrophy of mammalian skin, preventing, retarding, and/or treating the appearance of dark, under-eye circles and/or puffy eyes, preventing, retarding, and/or treating sallowness of mammalian skin, preventing and/or retarding tanning of mammalian skin, regulating and/or reducing the size of pores in mammalian skin, desquamating, exfoliating, and/or increasing turnover of mammalian skin, regulating oily/shiny appearance, preventing/retarding post-inflammatory hyperpigmentation in mammalian skin, and preventing and/or treating the appearance of cellulite in mammalian skin.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**METHODS FOR REGULATING THE CONDITION
OF MAMMALIAN KERATINOUS TISSUE VIA
TOPICAL APPLICATION OF PHYTOSTEROL COMPOSITIONS**

Cross Reference to Related Application

This application claims the benefit of U.S. Provisional Application No. 60/134,397, filed May 17, 1999.

Technical Field

The present invention relates to methods of regulating the condition of mammalian keratinous tissue using defined phytosterols wherein the methods include: a) thickening skin and preventing, retarding, and/or treating atrophy in mammalian skin, b) preventing, retarding, and/or treating the appearance of dark, under-eye circles and puffy eyes, c) preventing, retarding, and/or treating sallowness of mammalian skin, d) preventing and/or retarding tanning of mammalian skin, e) desquamating, exfoliating, and/or increasing turnover of mammalian skin, f) regulating and/or reducing the size of pores in mammalian skin, g) regulating the oily and/or shiny appearance of mammalian skin, h) preventing, retarding, and/or treating post-inflammatory hyperpigmentation, and i) preventing and/or treating the appearance of cellulite in mammalian skin. These methods are accomplished via the topical application of compositions containing specific phytosterols to the skin of a mammal needing such treatments.

Background of the Invention

Currently, there are a number of personal care products that are available to consumers, which are directed toward improving the health and physical appearance of keratinous tissues such as the skin, hair, and nails. The majority of these products are directed to delaying, minimizing or even eliminating skin wrinkling and other histological changes typically associated with the aging of skin or environmental damage to human skin.

Mammalian keratinous tissue, particularly human skin, is subjected to a variety of insults by both extrinsic and intrinsic factors. Such extrinsic factors include ultraviolet radiation, environmental pollution, wind, heat, infrared radiation, low humidity, harsh surfactants, abrasives, etc.. Intrinsic factors, on the other hand, include chronological aging and other biochemical changes from within the skin. Whether extrinsic or intrinsic, these factors result in visible signs of skin damage. Typical skin damage includes thinning of the skin, which occurs naturally as one ages. With such thinning, there is a reduction in the cells and blood vessels that supply the skin as well as a flattening of the dermal-epidermal junction that results in weaker mechanical resistance of this junction. See, for example, Oikarinen, "The Aging of Skin: Chronoaging Versus Photoaging," *Photodermatol. Photoimmunol. Photomed.*, vol. 7, pp. 3-4, 1990. Other damage or changes seen in aging or damaged skin includes sallowness, sagging, dark, under-eye circles, puffy eyes, enlarged pores, diminished rate of turnover, and abnormal desquamation or exfoliation.

Additional damage incurred as a result of both external and internal factors includes visible dead skin (i.e., flaking, scaling, dryness).

Therefore, there is a need for products and methods that seek to remedy these keratinous tissue conditions such that the conditions of keratinous tissues are regulated.

Without being limited by theory, it has been found that certain phytosterols can stabilize cellular and extracellular components of keratinous tissue, particularly the outer cell membranes as well as membranes of various organelles. The phytosterol treated cells become stronger, more resistant, and are able to withstand higher levels of the above-described insults.

Consequently, Applicants have surprisingly found that topical compositions that contain specific phytosterols may be used to provide prophylactic as well as therapeutic treatments for keratinous tissue conditions. For instance, Applicants have found that such compositions may be useful for treating atrophy of the skin as well as sallowness, spider vessels, dark, under-eye circles, puffy eyes, promoting skin desquamation, exfoliation, and/or turnover, regulating and/or reducing pore size appearance, preventing/retarding tanning, regulating oily/shiny appearance, preventing/retarding post-inflammatory hyperpigmentation in mammalian skin, and preventing and/or treating the appearance of cellulite in mammalian skin.

Summary of the Invention

The present invention relates to methods for regulating the condition of mammalian keratinous tissue wherein the methods each comprise the step of topically applying to the keratinous tissue of a mammal needing such treatment, a safe and effective amount of a composition comprising:

- a) a safe and effective amount of one or more phytosterols selected from the group consisting of β -sitosterol, campesterol, brassicasterol, $\Delta 5$ -avenasterol, lupenol, α -spinasterol, stigmasterol, their derivatives, and combinations thereof; and
- b) a dermatologically acceptable carrier for the phytosterol.

In additional embodiments, the above composition is suitable for thickening skin and preventing, retarding, and/or treating atrophy of mammalian skin, preventing, retarding, and/or treating the appearance of dark, under-eye circles and puffy eyes, preventing, retarding, and/or treating sallowness of mammalian skin, preventing and/or retarding tanning of mammalian skin, desquamating, exfoliating, or increasing the turnover of mammalian skin, regulating and/or reducing the size of pores in mammalian skin, regulating oily/shiny appearance of mammalian skin, preventing, retarding, and/or treating post-inflammatory hyperpigmentation, and preventing and/or treating the appearance of cellulite in mammalian skin.

Detailed Description of the Invention

All percentages and ratios used herein are by weight of the total composition and all measurements made are at 25°C, unless otherwise designated.

The compositions of the present invention can comprise, consist essentially of, or consist of, the essential components as well as optional ingredients described herein. As used herein, "consisting essentially of" means that the composition or component may include additional ingredients, but only if the

additional ingredients do not materially alter the basic and novel characteristics of the claimed compositions or methods.

All publications cited herein are hereby incorporated by reference in their entirety.

The term "keratinous tissue," as used herein, refers to keratin-containing layers disposed as the outermost protective covering of mammals which includes, but is not limited to, skin, hair, toenails, fingernails, cuticles, hooves, etc..

The term "topical application", as used herein, means to apply or spread the compositions of the present invention onto the surface of the keratinous tissue.

The term "dermatologically acceptable," as used herein, means that the compositions or components thereof so described are suitable for use in contact with human keratinous tissue without undue toxicity, incompatibility, instability, allergic response, and the like.

The term "safe and effective amount" as used herein means an amount of a compound or composition sufficient to significantly induce a positive benefit, preferably a positive keratinous tissue appearance or feel benefit, including independently or in combinations the benefits disclosed herein, but low enough to avoid serious side effects, i.e., to provide a reasonable benefit to risk ratio, within the scope of sound judgment of the skilled artisan.

The term "post-inflammatory hyperpigmentation" as used herein refers to the changes in melanin content as a response to an inflammatory event (e.g., acne, scratch, insect sting, sunburn, etc), especially in dark skin subjects.

The terms "desquamation, exfoliation, and/or increasing turnover" as used herein mean the removal of the upper layers of the stratum corneum (comprising the horny and granular layers). Without intending to be limited by theory, it is believed that these benefits may be accomplished via chemical and physical means that remove these layers from the top down. Additionally, it is possible to elicit exfoliation via a biological means that drives the turnover of the epidermal layers from the basal layers upwards. It is believed that this involves the process of keratinocyte proliferation as well as induction of differentiation. The latter leads to an elevation in keratinization levels as well, which ultimately lead to a reorganization of the upper epidermal layers that comprise the stratum corneum and stratum granular layers.

The terms "oily and/or shiny appearance" as used herein mean the glossy look mammalian skin tends to exhibit upon the excretion of oil, sebum, and/or sweat from the respective source gland.

The compositions of the present invention are useful for topical application and for regulating keratinous tissue condition. Regulation of keratinous tissue condition, especially human skin condition, is often required due to conditions that may be induced or caused by factors internal and/or external to the body. For instance, "regulating skin condition" includes prophylactically regulating and/or therapeutically regulating skin condition, and may involve one or more of the following benefits: thickening of skin (i.e., building the epidermis and/or dermis layers of the skin and/or the subcutaneous layers such as fat and muscle and where applicable the keratinous layers of the nail and hair shaft) to reduce skin atrophy, increasing the convolution of the dermal-epidermal border, non-melanin skin discoloration such as under eye circles,

blotching (e.g., uneven red coloration due to, e.g., rosacea) (hereinafter referred to as "red blotchiness"), sallowness (pale color), discoloration caused by telangiectasia or spider vessels. As used herein, prophylactically regulating skin condition includes delaying, minimizing and/or preventing visible and/or tactile discontinuities in skin (e.g., texture irregularities in the skin which may be detected visually or by feel). As used herein, therapeutically regulating skin condition includes ameliorating, e.g., diminishing, minimizing and/or effacing, discontinuities in skin. Regulating skin condition involves improving skin appearance and/or feel.

As used herein, "regulating skin condition" is intended to include regulation of such signs irrespective of the mechanism of origin.

The compositions of the present invention, including the essential and optional components thereof, are described in detail hereinafter.

Phytosterol

The topical compositions of the present invention comprise a safe and effective amount of one or more phytosterols selected from the group consisting of β -sitosterol, campesterol, brassicasterol, $\Delta 5$ -avenasterol, lupenol, α -spinasterol, stigmasterol, their derivatives, and combinations thereof. More preferably, the phytosterol is selected from the group consisting of β -sitosterol, campesterol, brassicasterol, stigmasterol, their derivatives, and combinations thereof. Even more preferably, the phytosterol is selected from the group consisting of β -sitosterol, campesterol, brassicasterol, stigmasterol, and combinations thereof. Most preferably, the phytosterol is stigmasterol.

Phytosterols can be synthetic or natural in origin and can be used as essentially pure compounds or mixtures of compounds (e.g., extracts from natural sources). Phytosterols are generally found in the unsaponifiable portion of vegetable oils and fats and are available as free sterols, acetylated derivatives, sterol esters, ethoxylated or glycosidic derivatives. More preferably, the phytosterols are free sterols. As used herein, "phytosterol" includes isomers and tautomers of such and is commercially available from Aldrich Chemical Company (Milwaukee, Wisconsin), Sigma Chemical Company (St. Louis, Missouri), and Fytokem Products, Inc. (Saskatoon, SK, Canada).

In the compositions of the present invention, the phytosterol preferably comprises from about 0.01% to about 99.99%, by weight of the composition, more preferably from about 0.01% to about 50%, even more preferably from about 0.1% to about 20%, still more preferably from about 0.2% to about 10%, and most preferably from about 0.5% to about 10%.

Dermatologically Acceptable Carrier

The topical compositions of the present invention also comprise a dermatologically acceptable carrier for the phytosterol. The phrase "dermatologically acceptable carrier", as used herein, means that the carrier is suitable for topical application to the keratinous tissue, has good aesthetic properties, is compatible with the actives of the present invention and any other components, and will not cause any safety or toxicity concerns. A safe and effective amount of carrier is from about 50% to about 99.99%,

preferably from about 80% to about 99.9%, more preferably from about 90% to about 98%, and most preferably from about 90% to about 95% of the composition.

The carrier can be in a wide variety of forms. For example, emulsion carriers, including, but not limited to, oil-in-water, water-in-oil, water-in-oil-in-water, and oil-in-water-in-silicone emulsions, are useful herein.

Preferred carriers comprise an emulsion such as oil-in-water emulsions and water-in-oil emulsions, e.g., silicone-in-water or water-in-silicone emulsions. As will be understood by the skilled artisan, a given component will distribute primarily into either the water or oil phase, depending on the water solubility/dispersability of the component in the composition. Phytosterol distributes primarily into the oil phase. Oil-in-water emulsions are especially preferred.

Emulsions according to the present invention generally contain a solution as described above and a lipid or oil. Lipids and oils may be derived from animals, plants, or petroleum and may be natural or synthetic (i.e., man-made). Preferred emulsions also contain a humectant, such as glycerin. Emulsions will preferably further contain from about 1% to about 10%, more preferably from about 2% to about 5%, of an emulsifier, based on the weight of the carrier. Emulsifiers may be nonionic, anionic or cationic. Suitable emulsifiers are disclosed in, for example, U.S. Patent 3,755,560, issued August 28, 1973, Dickert et al.; U.S. Patent 4,421,769, issued December 20, 1983, Dixon et al.; and McCutcheon's Detergents and Emulsifiers, North American Edition, pages 317-324 (1986).

The emulsion may also contain an anti-foaming agent to minimize foaming upon application to the keratinous tissue. Anti-foaming agents include high molecular weight silicones and other materials well known in the art for such use.

Suitable emulsions may have a wide range of viscosities, depending on the desired product form. Exemplary low viscosity emulsions, which are preferred, have a viscosity of about 50 centistokes or less, more preferably about 10 centistokes or less, most preferably about 5 centistokes or less.

Preferred water-in-silicone and oil-in-water emulsions are described in greater detail below.

a) Water-in-silicone emulsion

Water-in-silicone emulsions contain a continuous silicone phase and a dispersed aqueous phase.

(i) Continuous silicone phase

Preferred water-in-silicone emulsions of the present invention comprise from about 1% to about 60%, preferably from about 5% to about 40%, more preferably from about 10% to about 20%, by weight of a continuous silicone phase. The continuous silicone phase exists as an external phase that contains or surrounds the discontinuous aqueous phase described hereinafter.

The continuous silicone phase contains a polyorganosiloxane oil. A preferred water-in-silicone emulsion system is formulated to provide an oxidatively stable vehicle for an optional retinoid. The continuous silicone phase of these preferred emulsions comprises between about 50% and about 99.9% by weight of organopolysiloxane oil and less than about 50% by weight of a non-silicone oil. In a preferred embodiment, the continuous silicone phase comprises at least about 50%, preferably from about 60% to

about 99.9%, more preferably from about 70% to about 99.9%, and even more preferably from about 80% to about 99.9%, polyorganosiloxane oil by weight of the continuous silicone phase, and up to about 50% non-silicone oils, preferably less about 40%, more preferably less than about 30%, even more preferably less than about 10%, and most preferably less than about 2%, by weight of the continuous silicone phase. These preferred emulsion systems provide more oxidative stability to a retinoid over extended periods of time than comparable water-in-oil emulsions containing lower concentrations of the polyorganosiloxane oil. Concentrations of non-silicone oils in the continuous silicone phase are minimized or avoided altogether so as to further enhance oxidative stability of the selected retinoid in the compositions. Water-in-silicone emulsions of this type are described in copending U.S. Patent Application Serial No. 08/570,275, filed December 11, 1995, in the names of Joseph Michael Zukowski, Brent William Mason, Larry Richard Robinson and Greg George Hillebrand.

The organopolysiloxane oil for use in the composition may be volatile, non-volatile, or a mixture of volatile and non-volatile silicones. The term "nonvolatile" as used in this context refers to those silicones that are liquid under ambient conditions and have a flash point (under one atmospheric pressure) of or greater than about 100°C. The term "volatile" as used in this context refers to all other silicone oils. Suitable organopolysiloxanes can be selected from a wide variety of silicones spanning a broad range of volatilities and viscosities. Examples of suitable organopolysiloxane oils include polyalkylsiloxanes, cyclic polyalkylsiloxanes, and polyalkylarylsiloxanes.

Polyalkylsiloxanes useful in the composition herein include polyalkylsiloxanes with viscosities of from about 0.5 to about 1,000,000 centistokes at 25°C. Such polyalkylsiloxanes can be represented by the general chemical formula $R_3SiO[R_2SiO]_xSiR_3$ wherein R is an alkyl group having from one to about 30 carbon atoms (preferably R is methyl or ethyl, more preferably methyl; also mixed alkyl groups can be used in the same molecule), and x is an integer from 0 to about 10,000, chosen to achieve the desired molecular weight which can range to over about 10,000,000. Commercially available polyalkylsiloxanes include the polydimethylsiloxanes, which are also known as dimethicones, examples of which include the Vicasil[®] series sold by General Electric Company and the Dow Corning[®] 200 series sold by Dow Corning Corporation. Specific examples of suitable polydimethylsiloxanes include Dow Corning[®] 200 fluid having a viscosity of 0.65 centistokes and a boiling point of 100°C, Dow Corning[®] 225 fluid having a viscosity of 10 centistokes and a boiling point greater than 200°C, and Dow Corning[®] 200 fluids having viscosities of 50, 350, and 12,500 centistokes, respectively, and boiling points greater than 200°C. Suitable dimethicones include those represented by the chemical formula $(CH_3)_3SiO[(CH_3)_2SiO]_x[CH_3RSiO]_ySi(CH_3)_3$ wherein R is straight or branched chain alkyl having from two to about 30 carbon atoms and x and y are each integers of 1 or greater selected to achieve the desired molecular weight which can range to over about 10,000,000. Examples of these alkyl-substituted dimethicones include cetyl dimethicone and lauryl dimethicone.

Cyclic polyalkylsiloxanes suitable for use in the composition include those represented by the chemical formula $[\text{SiR}_2\text{-O}]_n$ wherein R is an alkyl group (preferably R is methyl or ethyl, more preferably methyl) and n is an integer from about 3 to about 8, more preferably n is an integer from about 3 to about 7, and most preferably n is an integer from about 4 to about 6. When R is methyl, these materials are typically referred to as cyclomethicones. Commercially available cyclomethicones include Dow Corning[®] 244 fluid having a viscosity of 2.5 centistokes, and a boiling point of 172°C, which primarily contains the cyclomethicone tetramer (i.e. n=4), Dow Corning[®] 344 fluid having a viscosity of 2.5 centistokes and a boiling point of 178°C, which primarily contains the cyclomethicone pentamer (i.e. n=5), Dow Corning[®] 245 fluid having a viscosity of 4.2 centistokes and a boiling point of 205°C, which primarily contains a mixture of the cyclomethicone tetramer and pentamer (i.e. n=4 and 5), and Dow Corning[®] 345 fluid having a viscosity of 4.5 centistokes and a boiling point of 217°, which primarily contains a mixture of the cyclomethicone tetramer, pentamer, and hexamer (i.e. n=4, 5, and 6).

Also useful are materials such as trimethylsiloxysilicate, which is a polymeric material corresponding to the general chemical formula $[(\text{CH}_3)_3\text{SiO}_{1/2}]_x[\text{SiO}_2]_y$, wherein x is an integer from about 1 to about 500 and y is an integer from about 1 to about 500. A commercially available trimethylsiloxysilicate is sold as a mixture with dimethicone as Dow Corning[®] 593 fluid.

Dimethiconols are also suitable for use in the composition. These compounds can be represented by the chemical formulas $\text{R}_3\text{SiO}[\text{R}_2\text{SiO}]_x\text{SiR}_2\text{OH}$ and $\text{HOR}_2\text{SiO}[\text{R}_2\text{SiO}]_x\text{SiR}_2\text{OH}$ wherein R is an alkyl group (preferably R is methyl or ethyl, more preferably methyl) and x is an integer from 0 to about 500, chosen to achieve the desired molecular weight. Commercially available dimethiconols are typically sold as mixtures with dimethicone or cyclomethicone (e.g. Dow Corning[®] 1401, 1402, and 1403 fluids).

Polyalkylaryl siloxanes are also suitable for use in the composition. Polymethylphenyl siloxanes having viscosities from about 15 to about 65 centistokes at 25°C are especially useful.

Preferred for use herein are organopolysiloxanes selected from the group consisting of polyalkylsiloxanes, alkyl substituted dimethicones, cyclomethicones, trimethylsiloxysilicates, dimethiconols, polyalkylaryl siloxanes, and mixtures thereof. More preferred for use herein are polyalkylsiloxanes and cyclomethicones. Preferred among the polyalkylsiloxanes are dimethicones.

As stated above, the continuous silicone phase may contain one or more non-silicone oils. Concentrations of non-silicone oils in the continuous silicone phase are preferably minimized or avoided altogether so as to further enhance oxidative stability of any retinoids in the compositions. Suitable non-silicone oils have a melting point of about 25°C or less under about one atmosphere of pressure. Examples of non-silicone oils suitable for use in the continuous silicone phase are those well known in the chemical arts in topical personal care products in the form of water-in-oil emulsions, e.g., mineral oil, vegetable oils, synthetic oils, semisynthetic oils, etc..

(ii) Dispersed aqueous phase

The topical compositions of the present invention comprise from about 30% to about 90%, more preferably from about 50% to about 85%, and most preferably from about 70% to about 80% of a dispersed aqueous phase. In emulsion technology, the term "dispersed phase" is a term well-known to one skilled in the art which means that the phase exists as small particles or droplets that are suspended in and surrounded by a continuous phase. The dispersed phase is also known as the internal or discontinuous phase. The dispersed aqueous phase is a dispersion of small aqueous particles or droplets suspended in and surrounded by the continuous silicone phase described hereinbefore.

The aqueous phase can be water, or a combination of water and one or more water soluble or dispersible ingredients. Nonlimiting examples of such optional ingredients include thickeners, acids, bases, salts, chelants, gums, water-soluble or dispersible alcohols and polyols, buffers, preservatives, sunscreens, agents, colorings, and the like.

The topical compositions of the present invention will typically comprise from about 25% to about 90%, preferably from about 40% to about 80%, more preferably from about 60% to about 80%, water in the dispersed aqueous phase by weight of the composition.

(iii) Emulsifier for dispersing the aqueous phase


The water-in-silicone emulsions of the present invention preferably comprise an emulsifier. In a preferred embodiment, the composition contains from about 0.1% to about 10% emulsifier, more preferably from about 0.5% to about 7.5%, most preferably from about 1% to about 5%, emulsifier by weight of the composition. The emulsifier helps disperse and suspend the aqueous phase within the continuous silicone phase.

A wide variety of emulsifying agents can be employed herein to form the preferred water-in-silicone emulsion. Known or conventional emulsifying agents can be used in the composition, provided that the selected emulsifying agent is chemically and physically compatible with essential components of the composition, and provides the desired dispersion characteristics. Suitable emulsifiers include silicone emulsifiers, non-silicon-containing emulsifiers, and mixtures thereof, known by those skilled in the art for use in topical personal care products. Preferably these emulsifiers have an HLB value of or less than about 14, more preferably from about 2 to about 14, and most preferably from about 4 to about 14. Emulsifiers having an HLB value outside of these ranges can be used in combination with other emulsifiers to achieve an effective weighted average HLB for the combination that falls within these ranges.

Silicone emulsifiers are preferred. A wide variety of silicone emulsifiers are useful herein. These silicone emulsifiers are typically organically modified organopolysiloxanes, also known to those skilled in the art as silicone surfactants. Useful silicone emulsifiers include dimethicone copolyols. These materials are polydimethylsiloxanes that have been modified to include polyether side chains such as polyethylene oxide chains, polypropylene oxide chains, mixtures of these chains, and polyether chains containing moieties derived from both ethylene oxide and propylene oxide. Other examples include alkyl-modified dimethicone copolyols, i.e., compounds that contain C2-C30 pendant side chains. Still other useful

dimethicone copolyols include materials having various cationic, anionic, amphoteric, and zwitterionic pendant moieties.


The dimethicone copolyol emulsifiers useful herein can be described by the following general structure:



wherein R is C1-C30 straight, branched, or cyclic alkyl and R² is selected from the group consisting of




and



wherein n is an integer from 3 to about 10; R³ and R⁴ are selected from the group consisting of H and C1-C6 straight or branched chain alkyl such that R³ and R⁴ are not simultaneously the same; and m, o, x, and y are selected such that the molecule has an overall molecular weight from about 200 to about 10,000,000, with m, o, x, and y being independently selected from integers of zero or greater such that m and o are not both simultaneously zero, and z being independently selected from integers of 1 or greater. It is recognized that positional isomers of these copolyols can be achieved. The chemical representations depicted above for the R² moieties containing the R³ and R⁴ groups are not meant to be limiting but are shown as such for convenience.

Also useful herein, although not strictly classified as dimethicone copolyols, are silicone surfactants as depicted in the structures in the previous paragraph wherein R² is:



wherein R⁵ is a cationic, anionic, amphoteric, or zwitterionic moiety.

Nonlimiting examples of dimethicone copolyols and other silicone surfactants useful as emulsifiers herein include polydimethylsiloxane polyether copolymers with pendant polyethylene oxide side chains, polydimethylsiloxane polyether copolymers with pendant polypropylene oxide side chains, polydimethylsiloxane polyether copolymers with pendant mixed polyethylene oxide and polypropylene oxide side chains, polydimethylsiloxane polyether copolymers with pendant mixed poly(ethylene)(propylene)oxide side chains, polydimethylsiloxane polyether copolymers with pendant organobetaine side chains, polydimethylsiloxane polyether copolymers with pendant carboxylate side chains, polydimethylsiloxane polyether copolymers with pendant quaternary ammonium side chains; and also further modifications of the preceding copolymers containing pendant C2-C30 straight, branched, or cyclic alkyl moieties. Examples of commercially available dimethicone copolyols useful herein sold by

Dow Corning Corporation are Dow Corning® 190, 193, Q2-5220, 2501 Wax, 2-5324 fluid, and 3225C (this latter material being sold as a mixture with cyclomethicone). Cetyl dimethicone copolyol is commercially available as a mixture with polyglyceryl-4 isostearate (and) hexyl laurate and is sold under the tradename ABIL® WE-09 (available from Goldschmidt). Cetyl dimethicone copolyol is also commercially available as a mixture with hexyl laurate (and) polyglyceryl-3 oleate (and) cetyl dimethicone and is sold under the tradename ABIL® WS-08 (also available from Goldschmidt). Other nonlimiting examples of dimethicone copolyols also include lauryl dimethicone copolyol, dimethicone copolyol acetate, diemethicone copolyol adipate, dimethicone copolyolamine, dimethicone copolyol behenate, dimethicone copolyol butyl ether, dimethicone copolyol hydroxy stearate, dimethicone copolyol isostearate, dimethicone copolyol laurate, dimethicone copolyol methyl ether, dimethicone copolyol phosphate, and dimethicone copolyol stearate. See International Cosmetic Ingredient Dictionary, Fifth Edition, 1993.

Dimethicone copolyol emulsifiers useful herein are described, for example, in U.S. Patent No. 4,960,764, to Figueroa, Jr. et al., issued October 2, 1990; European Patent No. EP 330,369, to Sanogueira, published August 30, 1989; G.H. Dahms, et al., "New Formulation Possibilities Offered by Silicone Copolyols," Cosmetics & Toiletries, vol. 110, pp. 91-100, March 1995; M.E. Carlotti et al., "Optimization of W/O-S Emulsions And Study Of The Quantitative Relationships Between Ester Structure And Emulsion Properties," J. Dispersion Science And Technology, 13(3), 315-336 (1992); P. Hameyer, "Comparative Technological Investigations of Organic and Organosilicone Emulsifiers in Cosmetic Water-in-Oil Emulsion Preparations," HAPPI 28(4), pp. 88-128 (1991); J. Smid-Korbar et al., "Efficiency and usability of silicone surfactants in emulsions," Provisional Communication, International Journal of Cosmetic Science, 12, 135-139 (1990); and D.G. Krzysik et al., "A New Silicone Emulsifier For Water-in-Oil Systems," Drug and Cosmetic Industry, vol. 146(4), pp. 28-81 (April 1990).

Among the non-silicone-containing emulsifiers useful herein are various non-ionic and anionic emulsifying agents such as sugar esters and polyesters, alkoxyated sugar esters and polyesters, C1-C30 fatty acid esters of C1-C30 fatty alcohols, alkoxyated derivatives of C1-C30 fatty acid esters of C1-C30 fatty alcohols, alkoxyated ethers of C1-C30 fatty alcohols, polyglyceryl esters of C1-C30 fatty acids, C1-C30 esters of polyols, C1-C30 ethers of polyols, alkyl phosphates, polyoxyalkylene fatty ether phosphates, fatty acid amides, acyl lactylates, soaps, and mixtures thereof. Other suitable emulsifiers are described, for example, in McCutcheon's, Detergents and Emulsifiers, North American Edition (1986), published by Allured Publishing Corporation; U.S. Patent No. 5,011,681 to Ciotti et al., issued April 30, 1991; U.S. Patent No. 4,421,769 to Dixon et al., issued December 20, 1983; and U.S. Patent No. 3,755,560 to Dickert et al., issued August 28, 1973.

Nonlimiting examples of these non-silicon-containing emulsifiers include: polyethylene glycol 20 sorbitan monolaurate (Polysorbate 20), polyethylene glycol 5 soya sterol, Steareth-20, Cetareth-20, PPG-2 methyl glucose ether distearate, Ceteth-10, Polysorbate 80, cetyl phosphate, potassium cetyl phosphate,

diethanolamine cetyl phosphate, Polysorbate 60, glyceryl stearate, PEG-100 stearate, polyoxyethylene 20 sorbitan trioleate (Polysorbate 85), sorbitan monolaurate, polyoxyethylene 4 lauryl ether sodium stearate, polyglyceryl-4 isostearate, hexyl laurate, steareth-20, cetareth-20, PPG-2 methyl glucose ether distearate, ceteth-10, diethanolamine cetyl phosphate, glyceryl stearate, PEG-100 stearate, and mixtures thereof.

b) Oil-in-Water Emulsions

Other preferred topical carriers include oil-in-water emulsions, having a continuous aqueous phase and a hydrophobic, water-insoluble phase ("oil phase") dispersed therein. Examples of suitable carriers comprising oil-in-water emulsions are described in U.S. Pat. No. 5,073,371, to Turner, D.J. et al., issued Dec. 17, 1991, and U.S. Pat. No. 5,073,372, to Turner, D.J. et al., issued Dec. 17, 1991. An especially preferred oil-in-water emulsion, containing a structuring agent, hydrophilic surfactant and water, is described in detail hereinafter.

(i) Structuring Agent

A preferred oil-in-water emulsion comprises a structuring agent to assist in the formation of a liquid crystalline gel network structure. Without being limited by theory, it is believed that the structuring agent assists in providing rheological characteristics to the composition that contribute to the stability of the composition. The structuring agent may also function as an emulsifier or surfactant. Preferred compositions of this invention comprise from about 0.5% to about 20%, more preferably from about 1% to about 10%, most preferably from about 1% to about 5%, by weight of the composition, of a structuring agent.

The preferred structuring agents of the present invention are selected from the group consisting of stearic acid, palmitic acid, stearyl alcohol, cetyl alcohol, behenyl alcohol, stearic acid, palmitic acid, the polyethylene glycol ether of stearyl alcohol having an average of about 1 to about 21 ethylene oxide units, the polyethylene glycol ether of cetyl alcohol having an average of about 1 to about 5 ethylene oxide units, and mixtures thereof. More preferred structuring agents of the present invention are selected from the group consisting of stearyl alcohol, cetyl alcohol, behenyl alcohol, the polyethylene glycol ether of stearyl alcohol having an average of about 2 ethylene oxide units (steareth-2), the polyethylene glycol ether of stearyl alcohol having an average of about 21 ethylene oxide units (steareth-21), the polyethylene glycol ether of cetyl alcohol having an average of about 2 ethylene oxide units, and mixtures thereof. Even more preferred structuring agents are selected from the group consisting of stearic acid, palmitic acid, stearyl alcohol, cetyl alcohol, behenyl alcohol, steareth-2, steareth-21, and mixtures thereof.

(ii) Hydrophilic surfactant


The preferred oil-in-water emulsions comprise from about 0.05% to about 10%, preferably from about 1% to about 6%, and more preferably from about 1% to about 3% of at least one hydrophilic surfactant which can disperse the hydrophobic materials in the water phase (percentages by weight of the topical carrier). The surfactant, at a minimum, must be hydrophilic enough to disperse in water.

Suitable surfactants include any of a wide variety of known cationic, anionic, zwitterionic, and amphoteric surfactants. See, McCutcheon's, Detergents and Emulsifiers, North American Edition (1986),

published by Allured Publishing Corporation; U.S. Patent 5,011,681; U.S. Patent 4,421,769; and U.S. Patent 3,755,560; these references are incorporated herein by reference in their entirety.

The exact surfactant chosen will depend upon the pH of the composition and the other components present.

Preferred are cationic surfactants, especially dialkyl quaternary ammonium compounds, examples of which are described in U.S. Patent 5,151,209; U.S. Patent 5,151,210; U.S. Patent 5,120,532; U.S. Patent 4,387,090; U.S. Patent 3,155,591; U.S. Patent 3,929,678; U.S. Patent 3,959,461; McCutcheon's, Detergents & Emulsifiers, (North American edition 1979) M.C. Publishing Co.; and Schwartz, et al., Surface Active Agents, Their Chemistry and Technology, New York: Interscience Publishers, 1949; which descriptions are incorporated herein by reference. The cationic surfactants useful herein include cationic ammonium salts such as those having the formula:



wherein R_1 , is an alkyl group having from about 12 to about 30 carbon atoms, or an aromatic, aryl or alkaryl group having from about 12 to about 30 carbon atoms; R_2 , R_3 , and R_4 are independently selected from hydrogen, an alkyl group having from about 1 to about 22 carbon atoms, or aromatic, aryl or alkaryl groups having from about 12 to about 22 carbon atoms; and X is any compatible anion, preferably selected from the group consisting of chloride, bromide, iodide, acetate, phosphate, nitrate, sulfate, methyl sulfate, ethyl sulfate, tosylate, lactate, citrate, glycolate, and mixtures thereof. Additionally, the alkyl groups of R_1 , R_2 , R_3 , and R_4 can also contain ester and/or ether linkages, or hydroxy or amino group substituents (e.g., the alkyl groups can contain polyethylene glycol and polypropylene glycol moieties).

More preferably, R_1 is an alkyl group having from about 12 to about 22 carbon atoms; R_2 is selected from H or an alkyl group having from about 1 to about 22 carbon atoms; R_3 and R_4 are independently selected from H or an alkyl group having from about 1 to about 3 carbon atoms; and X is as described previously.

Most preferably, R_1 is an alkyl group having from about 12 to about 22 carbon atoms; R_2 , R_3 , and R_4 are selected from H or an alkyl group having from about 1 to about 3 carbon atoms; and X is as described previously.

Alternatively, other useful cationic emulsifiers include amino-amides, wherein in the above structure R_1 is alternatively $R_5\text{CONH}-(\text{CH}_2)_n$, wherein R_5 is an alkyl group having from about 12 to about 22 carbon atoms, and n is an integer from about 2 to about 6, more preferably from about 2 to about 4, and most preferably from about 2 to about 3. Nonlimiting examples of these cationic emulsifiers include stearamidopropyl PG-dimonium chloride phosphate, behenamidopropyl PG dimonium chloride, stearamidopropyl ethyldimonium ethosulfate, stearamidopropyl dimethyl (myristyl acetate) ammonium

chloride, stearamidopropyl dimethyl cetearyl ammonium tosylate, stearamidopropyl dimethyl ammonium chloride, stearamidopropyl dimethyl ammonium lactate, and mixtures thereof. Especially preferred is behenamidopropyl PG dimonium chloride.

Nonlimiting examples of quaternary ammonium salt cationic surfactants include those selected from the group consisting of cetyl ammonium chloride, cetyl ammonium bromide, lauryl ammonium chloride, lauryl ammonium bromide, stearyl ammonium chloride, stearyl ammonium bromide, cetyl dimethyl ammonium chloride, cetyl dimethyl ammonium bromide, lauryl dimethyl ammonium chloride, lauryl dimethyl ammonium bromide, stearyl dimethyl ammonium chloride, stearyl dimethyl ammonium bromide, cetyl trimethyl ammonium chloride, cetyl trimethyl ammonium bromide, lauryl trimethyl ammonium chloride, lauryl trimethyl ammonium bromide, stearyl trimethyl ammonium chloride, stearyl trimethyl ammonium bromide, lauryl dimethyl ammonium chloride, stearyl dimethyl cetyl ditallow dimethyl ammonium chloride, dicetyl ammonium chloride, dicetyl ammonium bromide, dilauryl ammonium chloride, dilauryl ammonium bromide, distearyl ammonium chloride, distearyl ammonium bromide, dicetyl methyl ammonium chloride, dicetyl methyl ammonium bromide, dilauryl methyl ammonium chloride, dilauryl methyl ammonium bromide, distearyl methyl ammonium chloride, distearyl methyl ammonium bromide, and mixtures thereof. Additional quaternary ammonium salts include those wherein the C₁₂ to C₃₀ alkyl carbon chain is derived from a tallow fatty acid or from a coconut fatty acid. The term "tallow" refers to an alkyl group derived from tallow fatty acids (usually hydrogenated tallow fatty acids), which generally have mixtures of alkyl chains in the C₁₆ to C₁₈ range. The term "coconut" refers to an alkyl group derived from a coconut fatty acid, which generally have mixtures of alkyl chains in the C₁₂ to C₁₄ range. Examples of quaternary ammonium salts derived from these tallow and coconut sources include ditallow dimethyl ammonium chloride, ditallow dimethyl ammonium methyl sulfate, di(hydrogenated tallow) dimethyl ammonium chloride, di(hydrogenated tallow) dimethyl ammonium acetate, ditallow dipropyl ammonium phosphate, ditallow dimethyl ammonium nitrate, di(coconutalkyl)dimethyl ammonium chloride, di(coconutalkyl)dimethyl ammonium bromide, tallow ammonium chloride, coconut ammonium chloride, stearamidopropyl PG-dimonium chloride phosphate, stearamidopropyl ethyldimonium ethosulfate, stearamidopropyl dimethyl (myristyl acetate) ammonium chloride, stearamidopropyl dimethyl cetearyl ammonium tosylate, stearamidopropyl dimethyl ammonium chloride, stearamidopropyl dimethyl ammonium lactate, and mixtures thereof. An example of a quaternary ammonium compound having an alkyl group with an ester linkage is ditallowyl oxyethyl dimethyl ammonium chloride.

More preferred cationic surfactants are those selected from the group consisting of behenamidopropyl PG dimonium chloride, dilauryl dimethyl ammonium chloride, distearyl dimethyl ammonium chloride, dimyristyl dimethyl ammonium chloride, dipalmityl dimethyl ammonium chloride, distearyl dimethyl ammonium chloride, stearamidopropyl PG-dimonium chloride phosphate, stearamidopropyl ethyldiammonium ethosulfate, stearamidopropyl dimethyl (myristyl acetate) ammonium


chloride, stearamidopropyl dimethyl cetearyl ammonium tosylate, stearamidopropyl dimethyl ammonium chloride, stearamidopropyl dimethyl ammonium lactate, and mixtures thereof.

Most preferred cationic surfactants are those selected from the group consisting of behenamidopropyl PG dimonium chloride, dilauryl dimethyl ammonium chloride, distearyl dimethyl ammonium chloride, dimyristyl dimethyl ammonium chloride, dipalmityl dimethyl ammonium chloride, and mixtures thereof.

A preferred combination of cationic surfactant and structuring agent is behenamidopropyl PG dimonium chloride and/or behenyl alcohol, wherein the ratio is preferably optimized to maintained to enhance physical and chemical stability, especially when such a combination contains ionic and/or highly polar solvents. This combination is especially useful for delivery of sunscreens agents such as zinc oxide and octyl methoxycinnamate.

A wide variety of anionic surfactants are also useful herein. See, e.g., U.S. Patent No. 3,929,678, to Laughlin et al., issued December 30, 1975, which is incorporated herein by reference in its entirety. Nonlimiting examples of anionic surfactants include the alkoyl isethionates, and the alkyl and alkyl ether sulfates. The alkoyl isethionates typically have the formula $\text{RCO-OCH}_2\text{CH}_2\text{SO}_3\text{M}$ wherein R is alkyl or alkenyl of from about 10 to about 30 carbon atoms, and M is a water-soluble cation such as ammonium, sodium, potassium and triethanolamine. Nonlimiting examples of these isethionates include those alkoyl isethionates selected from the group consisting of ammonium cocoyl isethionate, sodium cocoyl isethionate, sodium lauroyl isethionate, sodium stearyl isethionate, and mixtures thereof.

The alkyl and alkyl ether sulfates typically have the respective formulae ROSO_3M and $\text{RO}(\text{C}_2\text{H}_4\text{O})_x\text{SO}_3\text{M}$, wherein R is alkyl or alkenyl of from about 10 to about 30 carbon atoms, x is from about 1 to about 10, and M is a water-soluble cation such as ammonium, sodium, potassium and triethanolamine. Another suitable class of anionic surfactants is the water-soluble salts of the organic, sulfuric acid reaction products of the general formula:



wherein R_1 is chosen from the group consisting of a straight or branched chain, saturated aliphatic hydrocarbon radical having from about 8 to about 24, preferably about 10 to about 16, carbon atoms; and M is a cation. Still other anionic synthetic surfactants include the class designated as succinamates, olefin sulfonates having about 12 to about 24 carbon atoms, and β -alkyloxy alkane sulfonates. Examples of these materials are sodium lauryl sulfate and ammonium lauryl sulfate.

Other anionic materials useful herein are soaps (i.e. alkali metal salts, e.g., sodium or potassium salts) of fatty acids, typically having from about 8 to about 24 carbon atoms, preferably from about 10 to about 20 carbon atoms. The fatty acids used in making the soaps can be obtained from natural sources such as, for instance, plant or animal-derived glycerides (e.g., palm oil, coconut oil, soybean oil, castor oil,

tallow, lard, etc.) The fatty acids can also be synthetically prepared. Soaps are described in more detail in U.S. Patent No. 4,557,853.

Amphoteric and zwitterionic surfactants are also useful herein. Examples of amphoteric and zwitterionic surfactants which can be used in the compositions of the present invention are those which are broadly described as derivatives of aliphatic secondary and tertiary amines in which the aliphatic radical can be straight or branched chain and wherein one of the aliphatic substituents contains from about 8 to about 22 carbon atoms (preferably C₈ - C₁₈) and one contains an anionic water solubilizing group, e.g., carboxy, sulfonate, sulfate, phosphate, or phosphonate. Examples are alkyl imino acetates, and iminodialkanoates and aminoalkanoates of the formulas $RN(CH_2)_mCO_2M$ and $RNH(CH_2)_mCO_2M$ wherein m is from 1 to 4, R is a C₈-C₂₂ alkyl or alkenyl, and M is H, alkali metal, alkaline earth metal ammonium, or alkanolammonium. Also included are imidazolinium and ammonium derivatives. Specific examples of suitable amphoteric surfactants include sodium 3-dodecyl-aminopropionate, sodium 3-dodecylaminopropane sulfonate, N-alkyltaurines such as the one prepared by reacting dodecylamine with sodium isethionate according to the teaching of U.S. Patent 2,658,072 which is incorporated herein by reference in its entirety; N-higher alkyl aspartic acids such as those produced according to the teaching of U.S. Patent 2,438,091 which is incorporated herein by reference in its entirety; and the products sold under the trade name "Miranol" and described in U.S. Patent 2,528,378, which is incorporated herein by reference in its entirety. Other examples of useful amphoterics include phosphates, such as coamidopropyl PG-dimonium chloride phosphate (commercially available as Monaquat PTC, from Mona Corp.).

Also useful herein as amphoteric or zwitterionic surfactants are the betaines. Examples of betaines include the higher alkyl betaines, such as coco dimethyl carboxymethyl betaine, lauryl dimethyl carboxymethyl betaine, lauryl dimethyl alphacarboxyethyl betaine, cetyl dimethyl carboxymethyl betaine, cetyl dimethyl betaine (available as Lonza 16SP from Lonza Corp.), lauryl bis-(2-hydroxyethyl) carboxymethyl betaine, stearyl bis-(2-hydroxypropyl) carboxymethyl betaine, oleyl dimethyl gamma-carboxypropyl betaine, lauryl bis-(2-hydroxypropyl)alpha-carboxyethyl betaine, coco dimethyl sulfopropyl betaine, stearyl dimethyl sulfopropyl betaine, lauryl dimethyl sulfoethyl betaine, lauryl bis-(2-hydroxyethyl) sulfopropyl betaine, and amidobetaines and amidosulfobetaines (wherein the $RCONH(CH_2)_3$ radical is attached to the nitrogen atom of the betaine), oleyl betaine (available as amphoteric Velvetex OLB-50 from Henkel), and cocamidopropyl betaine (available as Velvetex BK-35 and BA-35 from Henkel).

Other useful amphoteric and zwitterionic surfactants include the sultaines and hydroxysultaines such as cocamidopropyl hydroxysultaine (available as Mirataine CBS from Rhone-Poulenc), and the alkanoyl sarcosinates corresponding to the formula $RCON(CH_3)CH_2CH_2CO_2M$ wherein R is alkyl or alkenyl of about 10 to about 20 carbon atoms, and M is a water-soluble cation such as ammonium, sodium, potassium and trialkanolamine (e.g., triethanolamine), a preferred example of which is sodium lauroyl sarcosinate.

(iii) Water

The preferred oil-in-water emulsion comprises from about 25% to about 98%, preferably from about 65% to about 95%, more preferably from about 70% to about 90% water by weight of the topical carrier.

The hydrophobic phase is dispersed in the continuous aqueous phase. The hydrophobic phase may contain water insoluble or partially soluble materials such as are known in the art, including but not limited to the silicones described herein in reference to silicone-in-water emulsions, and other oils and lipids such as described above in reference to emulsions.

The topical compositions of the subject invention, including but not limited to lotions and creams, may comprise a dermatologically acceptable emollient. Such compositions preferably contain from about 2% to about 50% of the emollient. As used herein, "emollient" refers to a material useful for the prevention or relief of dryness, as well as for the protection of the skin. A wide variety of suitable emollients are known and may be used herein. Sagarin, Cosmetics, Science and Technology, 2nd Edition, Vol. 1, pp. 32-43 (1972), incorporated herein by reference, contains numerous examples of materials suitable as an emollient. A preferred emollient is glycerin. Glycerin is preferably used in an amount of from or about 0.001 to or about 20%, more preferably from or about 0.01 to or about 10%, most preferably from or about 0.1 to or about 5%, e.g., 3%.

Lotions and creams according to the present invention generally comprise a solution carrier system and one or more emollients. Lotions typically comprise from about 1% to about 20%, preferably from about 5% to about 10%, of emollient; from about 50% to about 90%, preferably from about 60% to about 80%, water; and phytosterol in the above described amounts. A cream typically comprises from about 5% to about 50%, preferably from about 10% to about 20%, of emollient; from about 45% to about 85%, preferably from about 50% to about 75%, water; and the phytosterol in the above described amounts.

Ointments of the present invention may comprise a simple carrier base of animal or vegetable oils or semi-solid hydrocarbons (oleaginous); absorption ointment bases which absorb water to form emulsions; or water soluble carriers, e.g., a water soluble solution carrier. Ointments may further comprise a thickening agent, such as described in Sagarin, Cosmetics, Science and Technology, 2nd Edition, Vol. 1, pp. 72-73 (1972), incorporated herein by reference, and/or an emollient. For example, an ointment may comprise from about 2% to about 10% of an emollient; from about 0.1% to about 2% of a thickening agent; and phytosterol in the above described amount.

Compositions of this invention useful for cleansing ("cleansers") are formulated with a suitable carrier, e.g., as described above, and preferably contain, in addition to the phytosterol in the above described amounts, from about 1% to about 90%, more preferably from about 5% to about 10%, of a dermatologically acceptable surfactant. The surfactant is suitably selected from anionic, nonionic, zwitterionic, amphoteric and ampholytic surfactants, as well as mixtures of these surfactants. Such surfactants are well known to those skilled in the detergency art. Nonlimiting examples of possible surfactants include isoceteth-20, sodium methyl cocoyl taurate, sodium methyl oleoyl taurate, and sodium

lauryl sulfate. See U.S. Patent No. 4,800,197, to Kowcz et al., issued January 24, 1989, which is incorporated herein by reference in its entirety, for exemplary surfactants useful herein. Examples of a broad variety of additional surfactants useful herein are described in McCutcheon's Detergents and Emulsifiers, North American Edition (1986), published by Allured Publishing Corporation. The cleansing compositions can optionally contain, at their art-established levels, other materials that are conventionally used in cleansing compositions.

The physical form of the cleansing compositions is not critical. The compositions can be, for example, formulated as toilet bars, liquids, shampoos, bath gels, hair conditioners, hair tonics, pastes, or mousses. Toilet bars are most preferred since this is the form of cleansing agent most commonly used to wash the skin. Rinse-off cleansing compositions, such as shampoos, require a delivery system adequate to deposit sufficient levels of actives on the skin and scalp. A preferred delivery system involves the use of insoluble complexes. For a more complete disclosure of such delivery systems, see U.S. Patent 4,835,148, Barford et al., issued May 30, 1989.

The compositions of the present invention may also be in the form of cosmetics. Suitable cosmetic forms include, but are not limited to, foundations, lipsticks, rouges, mascaras, and the like. Such cosmetic products may include conventional ingredients such as oils, colorants, pigments, emollients, fragrances, waxes, stabilizers, and the like. Exemplary carriers and such other ingredients which are suitable for use herein are described, for example, in copending patent application Serial No. 08/430,961, filed on April 28, 1995 in the names of Marcia L. Canter, Brain D. Barford, and Brian D. Hofrichter, and U.K. Patent Application GB 2274585-A, published on Jan. 23, 1993.

Optional Components

The compositions of the present invention may contain a variety of other ingredients such as are conventionally used in a given product type provided that they do not unacceptably alter the benefits of the invention.

In a preferred embodiment, where the composition is to be in contact with human keratinous tissue, the optional components should be suitable for application to keratinous tissue, that is, when incorporated into the composition they are suitable for use in contact with human keratinous tissue without undue toxicity, incompatibility, instability, allergic response, and the like within the scope of sound medical judgment. The *CTFA Cosmetic Ingredient Handbook*, Second Edition (1992) describes a wide variety of nonlimiting cosmetic and pharmaceutical ingredients commonly used in the skin care industry, which are suitable for use in the compositions of the present invention. Examples of these ingredient classes include: abrasives, absorbents, aesthetic components such as fragrances, pigments, colorings/colorants, essential oils, skin sensates, astringents, etc. (e.g., clove oil, menthol, camphor, eucalyptus oil, eugenol, menthyl lactate, witch hazel distillate), anti-acne agents, anti-caking agents, antifoaming agents, antimicrobial agents (e.g., iodopropyl butylcarbamate), antioxidants, binders, biological additives, buffering agents, bulking agents, chelating agents, chemical additives, colorants, cosmetic astringents, cosmetic biocides, denaturants, drug astringents, external analgesics, film formers or materials, e.g., polymers, for aiding the

film-forming properties and substantivity of the composition (e.g., copolymer of eicosene and vinyl pyrrolidone), opacifying agents, pH adjusters, propellants, reducing agents, sequestrants, skin bleaching and lightening agents (e.g., hydroquinone, kojic acid, ascorbic acid, magnesium ascorbyl phosphate, ascorbyl glucosamine), skin-conditioning agents (e.g., humectants, including miscellaneous and occlusive), skin soothing and/or healing agents (e.g., panthenol and derivatives (e.g., ethyl panthenol), aloe vera, pantothenic acid and its derivatives, allantoin, bisabolol, and dipotassium glycyrrhizinate), skin treating agents, thickeners, and vitamins and derivatives thereof.

In any embodiment of the present invention, however, the actives useful herein can be categorized by the benefit they provide or by their postulated mode of action. However, it is to be understood that the actives useful herein can in some instances provide more than one benefit or operate via more than one mode of action. Therefore, classifications herein are made for the sake of convenience and are not intended to limit the active to that particular application or applications listed.

Desquamation Actives

A safe and effective amount of a desquamation active may be added to the compositions of the present invention, more preferably from about 0.1% to about 10%, even more preferably from about 0.2% to about 5%, also preferably from about 0.5% to about 4%, by weight of the composition. Desquamation actives enhance the skin appearance benefits of the present invention. For example, the desquamation actives tend to improve the texture of the skin (e.g., smoothness). One desquamation system that is suitable for use herein comprises sulfhydryl compounds and zwitterionic surfactants and is described in copending application Serial No. 08/480,632, filed on June 7, 1995 in the name of Donald L. Bissett, corresponding to PCT Application No. U.S. 95/08136, filed 6/29/95. Another desquamation system that is suitable for use herein comprises salicylic acid and zwitterionic surfactants and is described in copending patent application Serial No. 08/554,944, filed on November 13, 1995 as a continuation of Serial No. 08/209,401, filed on March 9, 1994 in the name of Bissett, corresponding to PCT Application No. 94/12745, filed 11/4/94, published 5/18/95. Zwitterionic surfactants such as described in these applications are also useful as desquamatory agents herein, with cetyl betaine being particularly preferred.

Anti-Acne Actives

The compositions of the present invention may comprise a safe and effective amount of one or more anti-acne actives. Examples of useful anti-acne actives include resorcinol, sulfur, salicylic acid, erythromycin, zinc, etc. Further examples of suitable anti-acne actives are described in further detail in U. S. Patent No. 5,607,980, issued to McAtee et al, on March 4, 1997.

Anti-Wrinkle Actives/Anti-Atrophy Actives


The compositions of the present invention may further comprise a safe and effective amount of one or more anti-wrinkle actives or anti-atrophy actives. Exemplary anti-wrinkle/anti-atrophy actives suitable for use in the compositions of the present invention include sulfur-containing D and L amino acids and their derivatives and salts, particularly the N-acetyl derivatives, a preferred example of which is N-acetyl-L-cysteine; thiols, e.g. ethane thiol; hydroxy acids, phytic acid, lipoic acid; lysophosphatidic acid, skin peel

agents (e.g., phenol and the like), flavonoids (e.g., flavanones, chalcones, isoflavones, flavones, etc.), terpene alcohols (e.g., farnesol), vitamin B₃ compounds and retinoids which enhance the keratinous tissue appearance benefits of the present invention, especially in regulating keratinous tissue condition, e.g., skin condition.

a) Vitamin B₃ Compounds

The compositions of the present invention may comprise a safe and effective amount of a vitamin B₃ compound. Vitamin B₃ compounds are particularly useful for regulating skin condition as described in co-pending U. S. Application Serial No. 08/834,010, filed April 11, 1997 (corresponding to international publication WO 97/39733 A1, published October 30, 1997). When vitamin B₃ compounds are present in the compositions of the instant invention, the compositions preferably comprise from about 0.01% to about 50%, more preferably from about 0.1% to about 10%, even more preferably from about 0.5% to about 10%, and still more preferably from about 1% to about 5%, most preferably from about 2% to about 5%, by weight of the composition, of the vitamin B₃ compound.

As used herein, "vitamin B₃ compound" means a compound having the formula:



wherein R is - CONH₂ (i.e., niacinamide), - COOH (i.e., nicotinic acid) or - CH₂OH (i.e., nicotiny alcohol); derivatives thereof; and salts of any of the foregoing.

Exemplary derivatives of the foregoing vitamin B₃ compounds include nicotinic acid esters, including non-vasodilating esters of nicotinic acid (e.g., tocopheryl nicotinate), nicotiny amino acids, nicotiny alcohol esters of carboxylic acids, nicotinic acid N-oxide and niacinamide N-oxide.

Examples of suitable vitamin B₃ compounds are well known in the art and are commercially available from a number of sources, e.g., the Sigma Chemical Company (St. Louis, MO); ICN Biomedicals, Inc. (Irvin, CA) and Aldrich Chemical Company (Milwaukee, WI).

The vitamin compounds may be included as the substantially pure material, or as an extract obtained by suitable physical and/or chemical isolation from natural (e.g., plant) sources.

b) Retinoids

The compositions of the present invention may also comprise a retinoid. As used herein, "retinoid" includes all natural and/or synthetic analogs of Vitamin A or retinol-like compounds which possess the biological activity of Vitamin A in the skin as well as the geometric isomers and stereoisomers of these compounds. The retinoid is preferably retinol, retinol esters (e.g., C₂ - C₂₂ alkyl esters of retinol, including retinyl palmitate, retinyl acetate, retinyl propionate), retinal, and/or retinoic acid (including all-trans retinoic acid and/or 13-cis-retinoic acid), more preferably retinoids other than retinoic acid. These compounds are well known in the art and are commercially available from a number of sources, e.g., Sigma Chemical Company (St. Louis, MO), and Boehringer Mannheim (Indianapolis, IN). Other retinoids which are useful herein are described in U.S. Patent Nos. 4,677,120, issued Jun. 30, 1987 to Parish et al.; 4,885,311, issued Dec. 5, 1989 to Parish et al.; 5,049,584, issued Sep. 17, 1991 to Purcell et al.; 5,124,356, issued Jun. 23, 1992 to Purcell et al.; and Reissue 34,075, issued Sep. 22, 1992 to Purcell et al.. Other suitable retinoids are tocopheryl-retinoate [tocopherol ester of retinoic acid (trans- or cis-), adapalene {6-[3-(1-adamantyl)-4-methoxyphenyl]-2-naphthoic acid}, and tazarotene (ethyl 6-[2-(4,4-dimethylthiochroman-6-yl)-ethynyl]nicotinate). Preferred retinoids are retinol, retinyl palmitate, retinyl acetate, retinyl propionate, retinal and combinations thereof.

The retinoid may be included as the substantially pure material, or as an extract obtained by suitable physical and/or chemical isolation from natural (e.g., plant) sources. The retinoid is preferably substantially pure, more preferably essentially pure.

The compositions of this invention may contain a safe and effective amount of the retinoid, such that the resultant composition is safe and effective for regulating keratinous tissue condition, preferably for regulating visible and/or tactile discontinuities in skin, more preferably for regulating signs of skin aging, even more preferably for regulating visible and/or tactile discontinuities in skin texture associated with skin aging. The compositions preferably contain from or about 0.005% to or about 2%, more preferably 0.01% to or about 2%, retinoid. Retinol is most preferably used in an amount of from or about 0.01% to or about 0.15%; retinol esters are most preferably used in an amount of from or about 0.01% to or about 2% (e.g., about 1%); retinoic acids are most preferably used in an amount of from or about 0.01% to or about 0.25%; tocopheryl-retinoate, adapalene, and tazarotene are most preferably used in an amount of from or about 0.01% to or about 2%.

Where the compositions of the present invention contain both a retinoid and a Vitamin B₃ compound, the retinoid is preferably used in the above amounts, and the vitamin B₃ compound is preferably used in an amount of from or about 0.1% to or about 10%, more preferably from or about 2% to or about 5%.

Anti-Oxidants/Radical Scavengers

The compositions of the present invention may include a safe and effective amount of an anti-oxidant/radical scavenger. The anti-oxidant/radical scavenger is especially useful for providing protection

against UV radiation that can cause increased scaling or texture changes in the stratum corneum and against other environmental agents, which can cause skin damage.

A safe and effective amount of an anti-oxidant/radical scavenger may be added to the compositions of the subject invention, preferably from about 0.1% to about 10%, more preferably from about 1% to about 5%, of the composition.

Anti-oxidants/radical scavengers such as ascorbic acid (vitamin C) and its salts, ascorbyl esters of fatty acids, ascorbic acid derivatives (e.g., magnesium ascorbyl phosphate), tocopherol (vitamin E), tocopherol sorbate, tocopherol acetate, other esters of tocopherol, butylated hydroxy benzoic acids and their salts, 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (commercially available under the tradename Trolox^R), gallic acid and its alkyl esters, especially propyl gallate, uric acid and its salts and alkyl esters, sorbic acid and its salts, lipoic acid, amines (e.g., N,N-diethylhydroxylamine, amino-guanidine), sulfhydryl compounds (e.g., glutathione), dihydroxy fumaric acid and its salts, lysine pidolate, arginine pilolate, nordihydroguaiaretic acid, bioflavonoids, lysine, methionine, proline, superoxide dismutase, silymarin, tea extracts, grape skin/seed extracts, melanin, and rosemary extracts may be used. Preferred anti-oxidants/radical scavengers are selected from tocopherol sorbate and other esters of tocopherol, more preferably tocopherol sorbate. For example, the use of tocopherol sorbate in topical compositions and applicable to the present invention is described in U.S. Patent No. 4,847,071, issued on July 11, 1989 to Donald L. Bissett, Rodney D. Bush and Ranjit Chatterjee.

Chelators

The compositions of the present invention may also comprise a safe and effective amount of a chelator or chelating agent. As used herein, "chelator" or "chelating agent" means an active agent capable of removing a metal ion from a system by forming a complex so that the metal ion cannot readily participate in or catalyze chemical reactions. The inclusion of a chelating agent is especially useful for providing protection against UV radiation that can contribute to excessive scaling or skin texture changes and against other environmental agents, which can cause skin damage.

A safe and effective amount of a chelating agent may be added to the compositions of the subject invention, preferably from about 0.1% to about 10%, more preferably from about 1% to about 5%, of the composition. Exemplary chelators that are useful herein are disclosed in U.S. Patent No. 5,487,884, issued 1/30/96 to Bissett et al.; International Publication No. 91/16035, Bush et al., published 10/31/95; and International Publication No. 91/16034, Bush et al., published 10/31/95. Preferred chelators useful in compositions of the subject invention are furildioxime and derivatives thereof.

Flavonoids

The compositions of the present invention may optionally comprise a flavonoid compound. Flavonoids are broadly disclosed in U.S. Patents 5,686,082 and 5,686,367, both of which are herein incorporated by reference. Flavonoids suitable for use in the present invention are flavanones selected from the group consisting of unsubstituted flavanones, mono-substituted flavanones, and mixtures thereof; chalcones selected from the group consisting of unsubstituted chalcones, mono-substituted chalcones, di-

substituted chalcones, tri-substituted chalcones, and mixtures thereof; flavones selected from the group consisting of unsubstituted flavones, mono-substituted flavones, di-substituted flavones, and mixtures thereof; one or more isoflavones; coumarins selected from the group consisting of unsubstituted coumarins, mono-substituted coumarins, di-substituted coumarins, and mixtures thereof; chromones selected from the group consisting of unsubstituted chromones, mono-substituted chromones, di-substituted chromones, and mixtures thereof; one or more dicoumarols; one or more chromanones; one or more chromanols; isomers (e.g., cis/trans isomers) thereof; and mixtures thereof. By the term "substituted" as used herein means flavonoids wherein one or more hydrogen atom of the flavonoid has been independently replaced with hydroxyl, C1-C8 alkyl, C1-C4 alkoxy, O-glycoside, and the like or a mixture of these substituents.

Examples of suitable flavonoids include, but are not limited to, unsubstituted flavanone, mono-hydroxy flavanones (e.g., 2'-hydroxy flavanone, 6-hydroxy flavanone, 7-hydroxy flavanone, etc.), mono-alkoxy flavanones (e.g., 5-methoxy flavanone, 6-methoxy flavanone, 7-methoxy flavanone, 4'-methoxy flavanone, etc.), unsubstituted chalcone (especially unsubstituted trans-chalcone), mono-hydroxy chalcones (e.g., 2'-hydroxy chalcone, 4'-hydroxy chalcone, etc.), di-hydroxy chalcones (e.g., 2', 4'-dihydroxy chalcone, 2',4'-dihydroxy chalcone, 2,2'-dihydroxy chalcone, 2',3'-dihydroxy chalcone, 2',5'-dihydroxy chalcone, etc.), and tri-hydroxy chalcones (e.g., 2',3',4'-trihydroxy chalcone, 4,2',4'-trihydroxy chalcone, 2,2',4'-trihydroxy chalcone, etc.), unsubstituted flavone, 7,2'-dihydroxy flavone, 3',4'-dihydroxy naphthoflavone, 4'-hydroxy flavone, 5,6-benzoflavone, and 7,8-benzoflavone, unsubstituted isoflavone, daidzein (7,4'-dihydroxy isoflavone), 5,7-dihydroxy-4'-methoxy isoflavone, soy isoflavones (a mixture extracted from soy), unsubstituted coumarin, 4-hydroxy coumarin, 7-hydroxy coumarin, 6-hydroxy-4-methyl coumarin, unsubstituted chromone, 3-formyl chromone, 3-formyl-6-isopropyl chromone, unsubstituted dicoumarol, unsubstituted chromanone, unsubstituted chromanol, and mixtures thereof.

Preferred for use herein are unsubstituted flavanone, methoxy flavanones, unsubstituted chalcone, 2', 4'-dihydroxy chalcone, and mixtures thereof. Most preferred are unsubstituted flavanone, unsubstituted chalcone (especially the trans isomer), and mixtures thereof.

They can be synthetic materials or obtained as extracts from natural sources (e.g., plants). The naturally sourced material can also further be derivatized (e.g., an ester or ether derivative prepared following extraction from a natural source). Flavonoid compounds useful herein are commercially available from a number of sources, e.g., Indofine Chemical Company, Inc. (Somerville, New Jersey), Steraloids, Inc. (Wilton, New Hampshire), and Aldrich Chemical Company, Inc. (Milwaukee, Wisconsin).

Mixtures of the above flavonoid compounds may also be used.

The herein described flavonoid compounds are preferably present in the instant invention at concentrations of from about 0.01% to about 20%, more preferably from about 0.1% to about 10%, and most preferably from about 0.5% to about 5%.

Anti-Inflammatory Agents

A safe and effective amount of an anti-inflammatory agent may be added to the compositions of the present invention, preferably from about 0.1% to about 10%, more preferably from about 0.5% to about

5%, of the composition. The anti-inflammatory agent enhances the skin appearance benefits of the present invention, e.g., such agents contribute to a more uniform and acceptable skin tone or color. The exact amount of anti-inflammatory agent to be used in the compositions will depend on the particular anti-inflammatory agent utilized since such agents vary widely in potency.

Steroidal anti-inflammatory agents, including but not limited to, corticosteroids such as hydrocortisone, hydroxyltriamcinolone, alpha-methyl dexamethasone, dexamethasone-phosphate, beclomethasone dipropionates, clobetasol valerate, desonide, desoxymethasone, desoxycorticosterone acetate, dexamethasone, dichlorisone, diflorasone diacetate, difluocortolone valerate, fluadrenolone, fluclorolone acetonide, fludrocortisone, flumethasone pivalate, fluosinolone acetonide, fluocinonide, flucortine butylesters, fluocortolone, fluprednidene (fluprednylidene) acetate, flurandrenolone, halcinonide, hydrocortisone acetate, hydrocortisone butyrate, methylprednisolone, triamcinolone acetonide, cortisone, cortodoxone, flucetonide, fludrocortisone, difluorosone diacetate, fluradrenolone, fludrocortisone, difluorosone diacetate, fluradrenolone acetonide, medrysone, amcinafel, amcinafide, betamethasone and the balance of its esters, chloroprednisone, chlorprednisone acetate, clocortelone, clescinolone, dichlorisone, difluprednate, flucoronide, flunisolide, fluoromethalone, fluperolone, fluprednisolone, hydrocortisone valerate, hydrocortisone cyclopentylpropionate, hydrocortamate, meprednisone, paramethasone, prednisolone, prednisone, beclomethasone dipropionate, triamcinolone, and mixtures thereof may be used. The preferred steroidal anti-inflammatory for use is hydrocortisone.

A second class of anti-inflammatory agents, which is useful in the compositions, includes the nonsteroidal anti-inflammatory agents. The varieties of compounds encompassed by this group are well known to those skilled in the art. For detailed disclosure of the chemical structure, synthesis, side effects, etc. of non-steroidal anti-inflammatory agents, one may refer to standard texts, including Anti-inflammatory and Anti-Rheumatic Drugs, K. D. Rainsford, Vol. I-III, CRC Press, Boca Raton, (1985), and Anti-inflammatory Agents, Chemistry and Pharmacology, 1, R. A. Scherrer, et al., Academic Press, New York (1974).

Specific non-steroidal anti-inflammatory agents useful in the composition invention include, but are not limited to:

- 1) the oxicams, such as piroxicam, isoxicam, tenoxicam, sudoxicam, and CP-14, 304;
- 2) the salicylates, such as aspirin, disalcid, benorylate, trilisate, safapryn, solprin, diflunisal, and fendosal;
- 3) the acetic acid derivatives, such as diclofenac, fenclofenac, indomethacin, sulindac, tolmetin, isoxepac, furofenac, tiopinac, zidometacin, acematacin, fentiazac, zomepirac, clindanac, oxepinac, felbinac, and ketorolac;
- 4) the fenamates, such as mefenamic, meclofenamic, flufenamic, niflumic, and tolfenamic acids;

5) the propionic acid derivatives, such as ibuprofen, naproxen, benoxaprofen, flurbiprofen, ketoprofen, fenoprofen, fenbufen, indoprofen, piroprofen, carprofen, oxaprozin, pranoprofen, mioprofen, tiroxaprofen, suprofen, alminoprofen, and tiaprofenic; and

6) the pyrazoles, such as phenylbutazone, oxyphenbutazone, feprazone, azapropazone, and trimethazone.

Mixtures of these non-steroidal anti-inflammatory agents may also be employed, as well as the dermatologically acceptable salts and esters of these agents. For example, etofenamate, a flufenamic acid derivative, is particularly useful for topical application. Of the nonsteroidal anti-inflammatory agents, ibuprofen, naproxen, ketoprofen, flufenamic acid, etofenamate, aspirin, mefenamic acid, meclofenamic acid, piroxicam and felbinac are preferred; ibuprofen, naproxen, ketoprofen, etofenamate, aspirin and flufenamic acid are most preferred.

Finally, so-called "natural" anti-inflammatory agents are useful in methods of the present invention. Such agents may suitably be obtained as an extract by suitable physical and/or chemical isolation from natural sources (e.g., plants, fungi, by-products of microorganisms). For example, candelilla wax, alpha-bisabolol, aloe vera, Manjistha (extracted from plants in the genus Rubia, particularly Rubia Cordifolia), and Guggal (extracted from plants in the genus Commiphora, particularly Commiphora Mukul), kola extract, chamomile, red clover extract, and sea whip extract, may be used.

Additional anti-inflammatory agents useful herein include allantoin and compounds of the Licorice (the plant genus/species Glycyrrhiza glabra) family, including glycyrrhetic acid, glycyrrhizic acid, and derivatives thereof (e.g., salts and esters). Suitable salts of the foregoing compounds include metal and ammonium salts. Suitable esters include C₂ - C₂₄ saturated or unsaturated esters of the acids, preferably C₁₀ - C₂₄, more preferably C₁₆ - C₂₄. Specific examples of the foregoing include oil soluble licorice extract, the glycyrrhizic and glycyrrhetic acids themselves, monoammonium glycyrrhizinate, monopotassium glycyrrhizinate, dipotassium glycyrrhizinate, 1-beta-glycyrrhetic acid, stearyl glycyrrhethinate, and 3-stearyl-oxy-glycyrrhethinic acid, and disodium 3-succinyl-oxy-beta-glycyrrhethinate. Stearyl glycyrrhethinate is preferred.

The active component of these anti-inflammatory agents (e.g., biabolol, glycyrrhethinate esters) may also be obtained via extraction from natural sources or prepared synthetically.

Anti-Cellulite Agents

The compositions of the present invention may also comprise a safe and effective amount of an anti-cellulite agent. Suitable agents may include, but are not limited to, xanthine compounds (e.g., caffeine, theophylline, theobromine, and aminophylline).


Topical Anesthetics

The compositions of the present invention may also comprise a safe and effective amount of a topical anesthetic. Examples of topical anesthetic drugs include benzocaine, lidocaine, bupivacaine, chlorprocaine, dibucaine, etidocaine, mepivacaine, tetracaine, dyclonine, hexylcaine, procaine, cocaine, ketamine, pramoxine, phenol, and pharmaceutically acceptable salts thereof.

Tanning Actives

The compositions of the present invention may comprise a tanning active. When present, it is preferable that the compositions comprise from about 0.1% to about 20%, more preferably from about 2% to about 7%, and most preferably from about 3% to about 6%, by weight of the composition, of dihydroxyacetone as an artificial tanning active.

Dihydroxyacetone, which is also known as DHA or 1,3-dihydroxy-2-propanone, is a white to off-white, crystalline powder. This material can be represented by the chemical formula $C_3H_6O_3$ and the following chemical structure.



The compound can exist as a mixture of monomers and dimers, with the dimers predominating in the solid crystalline state. Upon heating or melting, the dimers break down to yield the monomers. This conversion of the dimeric form to the monomeric form also occurs in aqueous solution. Dihydroxyacetone is also known to be more stable at acidic pH values. See The Merck Index, Tenth Edition, entry 3167, p. 463 (1983), and "Dihydroxyacetone for Cosmetics", E. Merck Technical Bulletin, 03-304 110, 319 897, 180 588.

Skin Lightening Agents

The compositions of the present invention may comprise a skin lightening agent. When used, the compositions preferably comprise from about 0.1% to about 10%, more preferably from about 0.2% to about 5%, also preferably from about 0.5% to about 2%, by weight of the composition, of a skin lightening agent. Suitable skin lightening agents include those known in the art, including kojic acid, arbutin, ascorbic acid and derivatives thereof, e.g., magnesium ascorbyl phosphate or sodium ascorbyl phosphate or other salts of ascorbyl phosphate. Skin lightening agents suitable for use herein also include those described in copending patent application Serial No. 08/479,935, filed on June 7, 1995 in the name of Hillebrand, corresponding to PCT Application No. U.S. 95/07432, filed 6/12/95; and copending patent application

Serial No. 08/390,152, filed on February 24, 1995 in the names of Kalla L. Kvalnes, Mitchell A. DeLong, Barton J. Bradbury, Curtis B. Motley, and John D. Carter, corresponding to PCT Application No. U.S. 95/02809, filed 3/1/95, published 9/8/95.

Antimicrobial and Antifungal Actives

The compositions of the present invention may comprise an antimicrobial or antifungal active. Such actives are capable of destroying microbes, preventing the development of microbes or preventing the pathogenic action of microbes. A safe and effective amount of an antimicrobial or antifungal active may be added to the present compositions, preferably, from about 0.001% to about 10%, more preferably from about 0.01% to about 5%, and most preferably from about 0.05% to about 2%.

Examples of antimicrobial and antifungal actives include β -lactam drugs, quinolone drugs, ciprofloxacin, norfloxacin, tetracycline, erythromycin, amikacin, 2,4,4'-trichloro-2'-hydroxy diphenyl ether, 3,4,4'-trichlorobanilide, phenoxyethanol, phenoxy propanol, phenoxyisopropanol, doxycycline, capreomycin, chlorhexidine, chlortetracycline, oxytetracycline, clindamycin, ethambutol, hexamidine isethionate, metronidazole, pentamidine, gentamicin, kanamycin, lineomycin, methacycline, methenamine, minocycline, neomycin, netilmicin, paromomycin, streptomycin, tobramycin, miconazole, tetracycline hydrochloride, erythromycin, zinc erythromycin, erythromycin estolate, erythromycin stearate, amikacin sulfate, doxycycline hydrochloride, capreomycin sulfate, chlorhexidine gluconate, chlorhexidine hydrochloride, chlortetracycline hydrochloride, oxytetracycline hydrochloride, clindamycin hydrochloride, ethambutol hydrochloride, metronidazole hydrochloride, pentamidine hydrochloride, gentamicin sulfate, kanamycin sulfate, lineomycin hydrochloride, methacycline hydrochloride, methenamine hippurate, methenamine mandelate, minocycline hydrochloride, neomycin sulfate, netilmicin sulfate, paromomycin sulfate, streptomycin sulfate, tobramycin sulfate, miconazole hydrochloride, ketoconazole, amantadine hydrochloride, amantadine sulfate, octopirox, parachlorometa xlenol, nystatin, tolnaftate, zinc pyrithione and clotrimazole.

Preferred examples of actives useful herein include those selected from the group consisting of salicylic acid, benzoyl peroxide, 3-hydroxy benzoic acid, glycolic acid, lactic acid, 4-hydroxy benzoic acid, acetyl salicylic acid, 2-hydroxybutanoic acid, 2-hydroxypentanoic acid, 2-hydroxyhexanoic acid, cis-retinoic acid, trans-retinoic acid, retinol, phytic acid, N-acetyl-L-cysteine, lipoic acid, azelaic acid, arachidonic acid, benzoylperoxide, tetracycline, ibuprofen, naproxen, hydrocortisone, acetaminophen, resorcinol, phenoxyethanol, phenoxypropanol, phenoxyisopropanol, 2,4,4'-trichloro-2'-hydroxy diphenyl ether, 3,4,4'-trichlorocarbaniide, octopirox, lidocaine hydrochloride, clotrimazole, miconazole, ketoconazole, neocycin sulfate, and mixtures thereof.

Sunscreen Actives

Exposure to ultraviolet light can result in excessive scaling and texture changes of the stratum corneum. Therefore, the compositions of the subject invention may optionally contain a sunscreen active. As used herein, "sunscreen active" includes both sunscreen agents and physical sunblocks. Suitable sunscreen actives may be organic or inorganic.

A wide variety of conventional sunscreen actives are suitable for use herein. Sagarin, et al., at Chapter VIII, pages 189 et seq., of Cosmetics Science and Technology (1972), discloses numerous suitable actives. Specific suitable sunscreen actives include, for example: p-aminobenzoic acid, its salts and its derivatives (ethyl, isobutyl, glyceryl esters; p-dimethylaminobenzoic acid); anthranilates (i.e., o-aminobenzoates; methyl, menthyl, phenyl, benzyl, phenylethyl, linalyl, terpinyl, and cyclohexenyl esters); salicylates (amyl, phenyl, octyl, benzyl, menthyl, glyceryl, and di-pro-pyleneglycol esters); cinnamic acid derivatives (menthyl and benzyl esters, a-phenyl cinnamionitrile; butyl cinnamoyl pyruvate); dihydroxycinnamic acid derivatives (umbelliferone, methylumbelliferone, methylaceto-umbelliferone); trihydroxy-cinnamic acid derivatives (esculetin, methylesculetin, daphnetin, and the glucosides, esculin and daphnin); hydrocarbons (diphenylbutadiene, stilbene); dibenzalacetone and benzalacetophenone; naphtholsulfonates (sodium salts of 2-naphthol-3,6-disulfonic and of 2-naphthol-6,8-disulfonic acids); dihydroxynaphthoic acid and its salts; o- and p-hydroxybiphenyldisulfonates; coumarin derivatives (7-hydroxy, 7-methyl, 3-phenyl); diazoles (2-acetyl-3-bromindazole, phenyl benzoxazole, methyl naphthoxazole, various aryl benzothiazoles); quinine salts (bisulfate, sulfate, chloride, oleate, and tannate); quinoline derivatives (8-hydroxyquinoline salts, 2-phenylquinoline); hydroxy- or methoxy-substituted benzophenones; uric and violuric acids; tannic acid and its derivatives (e.g., hexaethylether); (butyl carboto) (6-propyl piperonyl) ether; hydroquinone; benzophenones (oxybenzene, sulisobenzene, dioxybenzene, benzoescorcinol, 2,2',4,4'-tetrahydroxybenzophenone, 2,2'-dihydroxy-4,4'-dimethoxybenzophenone, octabenzene; 4-isopropylidibenzoylmethane; butylmethoxydibenzoylmethane; octocrylene; octocrylene; [3-(4'-methylbenzylidene bornan-2-one); 4-isopropyl-di-benzoylmethane; zinc oxide and titanium dioxide.

Of these, 2-ethylhexyl-p-methoxycinnamate (commercially available as PARSOL MCX), 4,4'-t-butyl methoxydibenzoyl-methane (commercially available as PARSOL 1789), 2-hydroxy-4-methoxybenzophenone, octyldimethyl-p-aminobenzoic acid, digalloyltriolate, 2,2-dihydroxy-4-methoxybenzophenone, ethyl-4-(bis(hydroxy-propyl))aminobenzoate, 2-ethylhexyl-2-cyano-3,3-diphenylacrylate, 2-ethylhexyl-salicylate, glyceryl-p-aminobenzoate, 3,3,5-tri-methylcyclohexylsalicylate, methylanthranilate, p-dimethyl-aminobenzoic acid or aminobenzoate, 2-ethylhexyl-p-dimethyl-amino-benzoate, 2-phenylbenzimidazole-5-sulfonic acid, 2-(p-dimethylaminophenyl)-5-sulfonicbenzoxazoic acid, octocrylene, zinc oxide, titanium dioxide, and mixtures of these compounds, are preferred.

More preferred organic sunscreen actives useful in the compositions useful in the subject invention are 2-ethylhexyl-p-methoxycinnamate, butylmethoxydibenzoyl-methane, 2-hydroxy-4-methoxybenzo-phenone, 2-phenylbenzimidazole-5-sulfonic acid, octyldimethyl-p-aminobenzoic acid, octocrylene, zinc oxide, titanium dioxide, and mixtures thereof.

Also particularly useful in the compositions are sunscreen actives such as those disclosed in U.S. Patent No. 4,937,370 issued to Sabatelli on June 26, 1990, and U.S. Patent No. 4,999,186 issued to Sabatelli & Spirnak on March 12, 1991. The sunscreens disclosed therein have, in a single molecule, two distinct chromophore moieties, which exhibit different ultra-violet radiation absorption

spectra. One of the chromophore moieties absorbs predominantly in the UVB radiation range and the other absorbs strongly in the UVA radiation range.

Preferred members of this class of sunscreens are 4-N, N-(2-ethylhexyl) methyl-aminobenzoic acid ester of 2,4-dihydroxybenzophenone; N, N-di-(2-ethylhexyl)-4-aminobenzoic acid ester with 4-hydroxydibenzoylmethane; 4-N, N-(2-ethylhexyl) methyl-aminobenzoic acid ester with 4-hydroxydibenzoylmethane; 4-N, N-(2-ethylhexyl) methyl-aminobenzoic acid ester of 2-hydroxy-4-(2-hydroxyethoxy) benzophenone; 4-N, N-(2-ethylhexyl)-methylaminobenzoic acid ester of 4-(2-hydroxyethoxy) dibenzoylmethane; N,N-di-(2-ethylhexyl)-4-aminobenzoic acid ester of 2-hydroxy-4-(2-hydroxyethoxy)benzophenone; and N,N-di-(2-ethylhexyl)-4-aminobenzoic acid ester of 4-(2-hydroxyethoxy)dibenzoylmethane and mixtures thereof.

Especially preferred sunscreen actives include 4,4'-t-butylmethoxydibenzoylmethane, 2-ethylhexyl-p-methoxycinnamate, phenyl benzimidazole sulfonic acid, octocrylene, zinc oxide, and titanium dioxide, and mixtures thereof.

A safe and effective amount of the sunscreen active is used, typically from about 1% to about 20%, more typically from about 2% to about 10% by weight of the composition. Exact amounts will vary depending upon the sunscreen chosen and the desired Sun Protection Factor (SPF).

Conditioning Agents

The compositions of the present invention may comprise a conditioning agent selected from the group consisting of humectants, moisturizers, or skin conditioners. A variety of these materials can be employed and each can be present at a level of from about 0.01% to about 20%, more preferably from about 0.1% to about 10%, and most preferably from about 0.5% to about 7% by weight of the composition. These materials include, but are not limited to, guanidine; urea; glycolic acid and glycolate salts (e.g. ammonium and quaternary alkyl ammonium); salicylic acid; lactic acid and lactate salts (e.g., ammonium and quaternary alkyl ammonium); aloe vera in any of its variety of forms (e.g., aloe vera gel); polyhydroxy compounds such as sorbitol, mannitol, glycerol, hexanetriol, butanetriol, propylene glycol, butylene glycol, hexylene glycol and the like; polyethylene glycols; sugars (e.g., melibiose) and starches; sugar and starch derivatives (e.g., alkoxylated glucose, fructose, sucrose, etc.); hyaluronic acid; lactamide monoethanolamine; acetamide monoethanolamine; and mixtures thereof. Also useful herein are the propoxylated glycerols described in U. S. Patent No. 4,976,953, to Orr et al, issued December 11, 1990.

Also useful are various C₁-C₃₀ monoesters and polyesters of sugars and related materials. These esters are derived from a sugar or polyol moiety and one or more carboxylic acid moieties. Such ester materials are further described in, U. S. Patent No. 2,831,854, U. S. Patent No. 4,005,196, to Jandacek, issued January 25, 1977; U. S. Patent No. 4,005,195, to Jandacek, issued January 25, 1977, U. S. Patent No. 5,306,516, to Letton et al, issued April 26, 1994; U. S. Patent No. 5,306,515, to Letton et al, issued April 26, 1994; U. S. Patent No. 5,305,514, to Letton et al, issued April 26, 1994; U. S. Patent No. 4,797,300, to Jandacek et al, issued January 10, 1989; U. S. Patent No. 3,963,699, to Rizzi et al, issued

June 15, 1976; U. S. Patent No. 4,518,772, to Volpenhein, issued May 21, 1985; and U. S. Patent No. 4,517,360, to Volpenhein, issued May 21, 1985.

Preferably, the conditioning agent is selected from the group consisting of glycerol, urea, guanidine, sucrose polyester, and combinations thereof.

Thickening Agent (including thickeners and gelling agents)

The compositions of the present invention can comprise one or more thickening agents, preferably from about 0.1% to about 5%, more preferably from about 0.1% to about 3%, and most preferably from about 0.25% to about 2%, by weight of the composition.

Nonlimiting classes of thickening agents include those selected from the group consisting of:

a) Carboxylic Acid Polymers

These polymers are crosslinked compounds containing one or more monomers derived from acrylic acid, substituted acrylic acids, and salts and esters of these acrylic acids and the substituted acrylic acids, wherein the crosslinking agent contains two or more carbon-carbon double bonds and is derived from a polyhydric alcohol. Polymers useful in the present invention are more fully described in U. S. Patent No. 5,087,445, to Haffey et al, issued February 11, 1992; U. S. Patent No. 4,509,949, to Huang et al, issued April 5, 1985; U. S. Patent No. 2,798,053, to Brown, issued July 2, 1957; and in *CTFA International Cosmetic Ingredient Dictionary*, Fourth Edition, 1991, pp. 12 and 80.

Examples of commercially available carboxylic acid polymers useful herein include the carbomers, which are homopolymers of acrylic acid crosslinked with allyl ethers of sucrose or pentaerytritol. The carbomers are available as the Carbopol® 900 series from B.F. Goodrich (e.g., Carbopol® 954). In addition, other suitable carboxylic acid polymeric agents include copolymers of C₁₀₋₃₀ alkyl acrylates with one or more monomers of acrylic acid, methacrylic acid, or one of their short chain (i.e., C₁₋₄ alcohol) esters, wherein the crosslinking agent is an allyl ether of sucrose or pentaerytritol. These copolymers are known as acrylates/C₁₀₋₃₀ alkyl acrylate crosspolymers and are commercially available as Carbopol® 1342, Carbopol® 1382, Pemulen TR-1, and Pemulen TR-2, from B.F. Goodrich. In other words, examples of carboxylic acid polymer thickeners useful herein are those selected from the group consisting of carbomers, acrylates/C₁₀₋₃₀ alkyl acrylate crosspolymers, and mixtures thereof.

b) Crosslinked Polyacrylate Polymers

The compositions of the present invention can optionally comprise crosslinked polyacrylate polymers useful as thickeners or gelling agents including both cationic and nonionic polymers, with the cationics being generally preferred. Examples of useful crosslinked nonionic polyacrylate polymers and crosslinked cationic polyacrylate polymers are those described in U. S. Patent No. 5,100,660, to Hawe et al, issued March 31, 1992; U. S. Patent No. 4,849,484, to Heard, issued July 18, 1989; U. S. Patent No. 4,835,206, to Farrar et al, issued May 30, 1989; U.S. Patent No. 4,628,078 to Glover et al issued December 9, 1986; U.S. Patent No. 4,599,379 to Flesher et al issued July 8, 1986; and EP 228,868, to Farrar et al, published July 15, 1987.

c) Polyacrylamide Polymers

The compositions of the present invention can optionally comprise polyacrylamide polymers, especially nonionic polyacrylamide polymers including substituted branched or unbranched polymers. Most preferred among these polyacrylamide polymers is the nonionic polymer given the CTFA designation polyacrylamide and isoparaffin and laureth-7, available under the Tradename Sepigel 305 from Seppic Corporation (Fairfield, NJ).

Other polyacrylamide polymers useful herein include multi-block copolymers of acrylamides and substituted acrylamides with acrylic acids and substituted acrylic acids. Commercially available examples of these multi-block copolymers include Hypan SR150H, SS500V, SS500W, SSSA100H, from Lipo Chemicals, Inc., (Patterson, NJ).

d) Polysaccharides

A wide variety of polysaccharides are useful herein. "Polysaccharides" refer to gelling agents that contain a backbone of repeating sugar (i.e., carbohydrate) units. Nonlimiting examples of polysaccharide gelling agents include those selected from the group consisting of cellulose, carboxymethyl hydroxyethylcellulose, cellulose acetate propionate carboxylate, hydroxyethylcellulose, hydroxyethyl ethylcellulose, hydroxypropylcellulose, hydroxypropyl methylcellulose, methyl hydroxyethylcellulose, microcrystalline cellulose, sodium cellulose sulfate, and mixtures thereof. Also useful herein are the alkyl-substituted celluloses. In these polymers, the hydroxy groups of the cellulose polymer is hydroxyalkylated (preferably hydroxyethylated or hydroxypropylated) to form a hydroxyalkylated cellulose which is then further modified with a C₁₀-C₃₀ straight chain or branched chain alkyl group through an ether linkage. Typically these polymers are ethers of C₁₀-C₃₀ straight or branched chain alcohols with hydroxyalkylcelluloses. Examples of alkyl groups useful herein include those selected from the group consisting of stearyl, isostearyl, lauryl, myristyl, cetyl, isocetyl, cocoyl (i.e. alkyl groups derived from the alcohols of coconut oil), palmityl, oleyl, linoleyl, linolenyl, ricinoleyl, behenyl, and mixtures thereof. Preferred among the alkyl hydroxyalkyl cellulose ethers is the material given the CTFA designation cetyl hydroxyethylcellulose, which is the ether of cetyl alcohol and hydroxyethylcellulose. This material is sold under the tradename Natrosol[®] CS Plus from Aqualon Corporation (Wilmington, DE).

Other useful polysaccharides include scleroglucans comprising a linear chain of (1-3) linked glucose units with a (1-6) linked glucose every three units, a commercially available example of which is Clearogel[™] CS11 from Michel Mercier Products Inc. (Mountainside, NJ).

e) Gums

Other thickening and gelling agents useful herein include materials that are primarily derived from natural sources. Nonlimiting examples of these gelling agent gums include materials selected from the group consisting of acacia, agar, algin, alginic acid, ammonium alginate, amylopectin, calcium alginate, calcium carrageenan, carnitine, carrageenan, dextrin, gelatin, gellan gum, guar gum, guar hydroxypropyltrimonium chloride, hectorite, hyaluroinic acid, hydrated silica, hydroxypropyl chitosan, hydroxypropyl guar, karaya gum, kelp, locust bean gum, natto gum, potassium alginate, potassium

carrageenan, propylene glycol alginate, sclerotium gum, sodium carboxymethyl dextran, sodium carrageenan, tragacanth gum, xanthan gum, and mixtures thereof.

Preferred compositions of the present invention include a thickening agent selected from the group consisting of carboxylic acid polymers, crosslinked polyacrylate polymers, polyacrylamide polymers, and mixtures thereof, more preferably selected from the group consisting of carboxylic acid polymers, polyacrylamide polymers, and mixtures thereof.

Composition Preparation

The compositions of the present invention are generally prepared by conventional methods such as are known in the art of making topical compositions. Such methods typically involve mixing of the ingredients in one or more steps to a relatively uniform state, with or without heating, cooling, application of vacuum, and the like.

Methods for Regulating Keratinous Tissue Condition

The compositions of the present invention are useful for regulating a number of mammalian keratinous tissue conditions. Such regulation of keratinous tissue conditions includes prophylactic and therapeutic regulation. More specifically, such regulating methods are directed to, but are not limited to, thickening keratinous tissue (i.e., building the epidermis and/or dermis and/or subcutaneous layers of the skin and where applicable the keratinous layers of the nail and hair shaft) and preventing, retarding, and/or treating atrophy of mammalian skin, preventing, retarding, and/or treating the appearance of dark, under-eye circles and/or puffy eyes, preventing, retarding, and/or treating sallowness of mammalian skin, preventing and/or retarding tanning of mammalian skin, desquamating, exfoliating, and/or increasing turnover in mammalian skin, reducing the size of pores in mammalian skin, regulating oily/shiny appearance of mammalian skin, preventing, retarding, and/or treating post-inflammatory hyperpigmentation, and preventing and/or treating the appearance of cellulite in mammalian skin.

Regulating keratinous tissue condition involves topically applying to the keratinous tissue a safe and effective amount of a composition of the present invention. The amount of the composition that is applied, the frequency of application and the period of use will vary widely depending upon the level of phytosterol and/or other components of a given composition and the level of regulation desired, e.g., in light of the level of keratinous tissue damage present or expected to occur.

In a preferred embodiment, the composition is chronically applied to the skin. By "chronic topical application" is meant continued topical application of the composition over an extended period during the subject's lifetime, preferably for a period of at least about one week, more preferably for a period of at least about one month, even more preferably for at least about three months, even more preferably for at least about six months, and more preferably still for at least about one year. While benefits are obtainable after various maximum periods of use (e.g., five, ten or twenty years), it is preferred that chronic applications continue throughout the subject's lifetime. Typically applications would be on the order of about once per day over such extended periods, however application rates can vary from about once per week up to about three times per day or more.